

References



A

- Abel, D. C., & Grubbs, D. R. (2020). *Shark biology and conservation: Essentials for educators, students and enthusiasts*. John Hopkins University Press.
- Abrantes, K. G., & Barnett, A. (2011). Intrapopulation variations in diet and habitat use in a marine apex predator, the broadnose sevengill shark *Notorynchus cepedianus*. *Marine Ecology Progress Series*, 442, 133–148. <https://doi.org/10.3354/meps09395>
- Ackerman, J. T., Kondratieff, M. C., Matern, S. A., & Cech, J. J. (2000). Tidal influence on spatial dynamics of leopard sharks, *Triakis semifasciata*, in Tomales Bay, California. *Environmental Biology of Fishes*, 58(1), 33–43. <https://doi.org/10.1023/A:1007657019696>
- Adenle, A. A., Stevens, C., & Bridgewater, P. (2015). Global conservation and management of biodiversity in developing countries: An opportunity for a new approach. *Environmental Science and Policy*, 45(2015), 104–108. <https://doi.org/10.1016/j.envsci.2014.10.002>
- Adkins, M. E., Simpfendorfer, C. A., & Tobin, A. J. (2016). Large tropical fishes and their use of the nearshore littoral, intertidal and subtidal habitat mosaic. *Marine and Freshwater Research*, 67(10), 1534–1545. <https://doi.org/10.1071/MF14339>
- Ajemian, M. J., Dolan, D., Graham, W. M., & Powers, S. P. (2011). First evidence of elasmobranch predation by a waterbird: Stingray attack and consumption by the Great Blue Heron (*Ardea herodias*). *Waterbirds*, 34(1), 117–120. <https://doi.org/10.1675/063.034.0116>
- Ajemian, M. J., & Powers, S. P. (2012). Habitat-specific feeding by cownose rays (*Rhinoptera bonasus*) of the northern Gulf of Mexico. *Environmental Biology of Fishes*, 95, 79–97.
- Ajemian, M. J., Powers, S. P., & Murdoch, T. J. T. (2012). Estimating the potential impacts of large mesopredators on benthic resources: Integrative assessment of spotted eagle ray foraging ecology in Bermuda. *PLoS ONE*, 7(7). <https://doi.org/10.1371/journal.pone.0040227>
- Alder, J., & Sumaila, U. R. (2004). Western Africa: A fish basket of Europe past and present. *J. Environ. Dev*, 13, 156–178. <https://doi.org/10.1177/1070496504266092>.
- Almojil, D. (2021). Local ecological knowledge of fisheries charts decline of sharks in data-poor regions. *Marine Policy*, 132(June), 104638. <https://doi.org/10.1016/j.marpol.2021.104638>
- Alongi, D. M. (2014). Carbon cycling and storage in mangrove forests. *Annual Review of Marine Science*, 6, 195–219.
- Alongi, D. M. (2018). *Blue carbon* (1st ed.). Springer International Publishing.
- Alves, J. A., Coelho, A. P., Carneiro, C., Nightingale, J., Barros, A. R., & Rocha, A. D. (2021). Ligações migratórias estabelecidas pelas aves limícolas do Arquipélago Dos Bijagós. *Lucanus – Revista de Ambiente e Sociedade*, V, 166–185.
- Amandè, M. J., Ariz, J., Chassot, E., Molina, A. D., Gaertner, D., Murua, H., & Chavance, P. (2010). Bycatch of the European purse seine tuna fishery in the Atlantic Ocean for the 2003–2007 period. *Aquatic Living Resources*, 23(4), 353–362. <https://doi.org/10.1051/alr/2011003>
- Ameyaw, G. A., Tsamenyi, M., McIlgorm, A., & Aheto, D. W. (2021). Challenges in the management of small-scale marine fisheries conflicts in Ghana. *Ocean and Coastal Management*, 211(July 2020), 105791. <https://doi.org/10.1016/j.ocecoaman.2021.105791>
- Anadón, J. D., Giménez, A., Ballestar, R., & Pérez, I. (2009). Evaluation of local ecological knowledge as a method for collecting extensive data on animal abundance. *Conservation Biology*, 23(3), 617–625. <https://doi.org/10.1111/j.1523-1739.2008.01145.x>

- Anderson, J. L., Anderson, C. M., Chu, J., Meredith, J., Asche, F., Sylvia, G., & Valderrama, D. (2015). The fishery performance indicators: A management tool for triple bottom line outcomes. *PLoS ONE*, 10(5), 1–20. <https://doi.org/10.1371/journal.pone.0122809>
- Anderson, S. C., Flemming, J. M., Watson, R., & Lotze, H. K. (2011). Rapid global expansion of invertebrate fisheries: Trends, drivers, and ecosystem effects. *PLoS ONE*, 6(3), 1–9. <https://doi.org/10.1371/journal.pone.0014735>
- Arendt, M. D., Olney, J. E., & Lucy, J. A. (2001). Stomach content analysis of cobia, *Rachycentron canadum*, from lower Chesapeake Bay. *Fishery Bulletin*, 99(4), 665–670.
- Arístegui, J., Barton, E. D., Álvarez-Salgado, X. A., Santos, A. M. P., Figueiras, F. G., Kifani, S., & Demarcq, H. (2009). Sub-regional ecosystem variability in the Canary Current upwelling. *Progress in Oceanography*, 83(1–4), 33–48. <https://doi.org/10.1016/j.pocean.2009.07.031>
- Assaneo, F., Coutinho, R. M., Lin, Y., Mantilla, C., & Lutscher, F. (2013). Dynamics and coexistence in a system with intraguild mutualism. *Ecological Complexity*, 14, 64–74. <https://doi.org/10.1016/j.ecocom.2012.10.004>
- Assessment, M. E. (2005). *Ecosystems and human well-being: Synthesis*. Island Press.
- Atwood, T. B., Connolly, R. M., Ritchie, E. G., Lovelock, C. E., Heithaus, M. R., Hays, G. C., & Macreadie, P. I. (2015). Predators help protect carbon stocks in blue carbon ecosystems. *Nature Climate Change*, 5(12), 1038–1045. <https://doi.org/10.1038/nclimate2763>
- Azzurro, E., Moschella, P., & Maynou, F. (2011). Tracking signals of change in Mediterranean fish diversity based on local ecological knowledge. *PLoS ONE*, 6(9), 1–8. <https://doi.org/10.1371/journal.pone.0024885>

B

- Bakker, J., Wangensteen, O. S., Chapman, D. D., Boussarie, G., Buddo, D., Guttridge, T. L., & Mariani, S. (2017). Environmental DNA reveals tropical shark diversity in contrasting levels of anthropogenic impact. *Scientific Reports*, 1–11. <https://doi.org/10.1038/s41598-017-17150-2>
- Ballantyne, J. S., & Robinson, J. W. (2010). Freshwater elasmobranchs: A review of their physiology and biochemistry. *Journal of Comparative Physiology B: Biochemical, Systemic, and Environmental Physiology*, 180(4), 475–493. <https://doi.org/10.1007/s00360-010-0447-0>
- Balmford, A., Gravestock, P., Hockley, N., McClean, C. J., & Roberts, C. M. (2004). The worldwide costs of marine protected areas. *Proceedings of the National Academy of Sciences*, 101(26), 9694–9697. <https://doi.org/10.1073/pnas.0403239101>
- Bandimere, A., Brenner, H., Casale, P., Dimatteo, A., Hurley, B., Hutchinson, B., & Wallace, B. (2021). Important marine turtle areas: Guidelines 1.0 (pp. 1–12). IUCN-SSC Marine Turtle Specialist Group.
- Bangley, C. W., Paramore, L., Shiffman, D. S., & Rulifson, R. A. (2018). Increased abundance and nursery habitat use of the bull shark (*Carcharhinus leucas*) in response to a changing environment in a warm-temperate estuary. *Scientific Reports*, 8(1), 6018. <https://doi.org/10.1038/s41598-018-24510-z>
- Barbier, E. B., Hacker, S. D., Kennedy, C., Koch, E. W., Stier, A. C., & Silliman, B. R. (2011). The value of estuarine and coastal ecosystem services. *Ecological Monographs*, 81(2), 169–193.
- Barker, M. J., & Schluessel, V. (2005). Managing global shark fisheries: Suggestions for prioritizing management strategies. *Aquatic Conservation: Marine and Freshwater Ecosystems*, 15(4), 325–347. <https://doi.org/10.1002/aqc.660>

- Barley, S. C., Clark, T. D., & Meeuwig, J. J. (2020). Ecological redundancy between coral reef sharks and predatory teleosts. *Reviews in Fish Biology and Fisheries*, 30(1), 153–172. <https://doi.org/10.1007/s11160-019-09588-6>
- Barley, S. C., Meekan, M. G., & Meeuwig, J. J. (2017). Species diversity, abundance, biomass, size and trophic structure of fish assemblages on coral reefs in relation to shark abundance. *Mar.* <https://doi.org/10.6084/m9.figshare.4009686>
- Barnes, M. A., & Turner, C. R. (2016). The ecology of environmental DNA and implications for conservation genetics. *Conservation Genetics*, 17(1), 1–17. <https://doi.org/10.1007/s10592-015-0775-4>
- Barnett, A., Abrantes, K., Stevens, J. D., Yick, J. L., Frusher, S. D., & Semmens, J. M. (2010). Predator-prey relationships and foraging ecology of a marine apex predator with a wide temperate distribution. *Marine Ecology Progress Series*, 416, 189–200. <https://doi.org/10.3354/meps08778>
- Barnett, A., Abrantes Ká, K. G., Seymour, J., & Fitzpatrick, R. (2012). Residency and spatial use by reef sharks of an isolated seamount and its implications for conservation. *PLoS ONE*, 7(5), 1–12. <https://doi.org/10.1371/journal.pone.0036574>
- Barnett, A., Braccini, M., Dudgeon, C. L., Payne, N. L., Abrantes, K. G., Sheaves, M., & Snelling, E. P. (2017). The utility of bioenergetics modelling in quantifying predation rates of marine apex predators: Ecological and fisheries implications. *Scientific Reports*, 7(1), 1–10. <https://doi.org/10.1038/s41598-017-13388-y>
- Barr, L. M., & Possingham, H. P. (2013). Are outcomes matching policy commitments in Australian marine conservation planning? *Marine Policy*, 42(2013), 39–48. <https://doi.org/10.1016/j.marpol.2013.01.012>
- Barría, C., Coll, M., & Navarro, J. (2015). Unravelling the ecological role and trophic relationships of uncommon and threatened elasmobranchs in the western Mediterranean Sea. *Marine Ecology Progress Series*, 539, 225–240. <https://doi.org/10.3354/meps11494>
- Barrios-O'Neill, D., Bertolini, C., & Collins, P. C. (2017). Trophic cascades and the transient keystone concept. *Biological Conservation*, 212(May), 191–195. <https://doi.org/10.1016/j.biocon.2017.06.011>
- Barros, N. B., & Wells, R. S. (1998). Prey and feeding patterns of resident bottlenose dolphins (*Tursiops truncatus*) in Sarasota Bay, Florida. *Journal of Mammalogy*, 79(3), 1045–1059. <https://doi.org/10.2307/1383114>
- Bartón, K. (2022). Package “MuMIn” multi-model inference [Computer software].
- Bascompte, J., Melián, C. J., & Sala, E. (2005). Interaction strength combinations and the overfishing of a marine food web. *Proceedings of the National Academy of Sciences of the United States of America*, 102(15), 5443–5447. <https://doi.org/10.1073/pnas.0501562102>
- Battley, P. F., Warnock, N., Tibbitts, T. L., Gill, R. E., Piersma, T., Hassell, C. J., & Riegen, A. C. (2012). Contrasting extreme long-distance migration patterns in bar-tailed godwits *Limosa lapponica*. *Journal of Avian Biology*, 43(1), 21–32. <https://doi.org/10.1111/j.1600-048X.2011.05473.x>
- Baum, J. K., & Myers, R. A. (2004). Shifting baselines and the decline of pelagic sharks in the Gulf of Mexico. *Ecology Letters*, 7, 135–145.
- Baum, J. K., Myers, R. A., Kehler, D. G., Worm, B., Harley, S. J., & Doherty, P. A. (2003). Collapse and conservation of shark populations in the Northwest Atlantic. In *Science* (Vol. 299, pp. 389–392). <https://doi.org/10.1126/science.1079777>

- Baum, J. K., & Worm, B. (2009). Cascading top-down effects of changing oceanic predator abundances. *Journal of Animal Ecology*, 78, 699–714. <https://doi.org/10.1111/j.1365-2656.2009.01531.x>.
- Beaudreau, A. H., & Levin, P. S. (2014). Advancing the use of local ecological knowledge for assessing data-poor species in coastal ecosystems. *Ecological Applications*, 24(2), 244–256. <https://doi.org/10.1890/13-0817.1>
- Begossi, A. (2010). Small-scale fisheries in Latin America: Management models and challenges. *Mast*, 9(2), 7–31.
- Belhabib, D., Cheung, W. W. L., Kroodsmas, D., Lam, V. W. Y., Underwood, P. J., & Virdin, J. (2019). Catching industrial fishing incursions into inshore waters of Africa from space. *Fish and Fisheries*, 379–392. <https://doi.org/10.1111/faf.12436>
- Belhabib, D., Gascuel, D., Abou Kane, E., Harper, S., Zeller, D., & Pauly, D. (2013). Preliminary estimation of realistic fisheries removals from Mauritania, 1950-2010. *Fisheries Centre Research Reports*, 20(3), 61–78.
- Belhabib, D., Koutob, V., Sall, A., Lam, V. W. Y., & Pauly, D. (2014). Fisheries catch misreporting and its implications: The case of Senegal. *Fisheries Research*, 151, 1–11. <https://doi.org/10.1016/j.fishres.2013.12.006>
- Belhabib, D., & Pauly, D. (2015). Fisheries in troubled waters: A catch reconstruction for Guinea-Bissau, 1950-2010. *Fisheries Centre Research Reports*, 23(3), 1–16.
- Bender, M. G., Machado, G. R., Azevedo Silva, P. J., Floeter, S. R., Monteiro-Netto, C., Luiz, O. J., & Ferreira, C. E. L. (2014). Local ecological knowledge and scientific data reveal overexploitation by multigear artisanal fisheries in the Southwestern Atlantic. *PLoS ONE*, 9(10). <https://doi.org/10.1371/journal.pone.0110332>
- Beninger, P. G. (2019). *Mudflat Ecology*. Aquatic Ecology Series, 7.
- Bennie, J. J., Duffy, J. P., Inger, R., & Gaston, K. J. (2014). Biogeography of time partitioning in mammals. *Proceedings of the National Academy of Sciences of the United States of America*, 111(38), 13727–13732. <https://doi.org/10.1073/pnas.1216063110>
- Bernal, D., Carlson, J. K., Goldman, K. J., & Lowe, C. G. (2012). Energetics, metabolism, and endothermy in sharks and rays. *Biology of Sharks and Their Relatives*, 227–254. <https://doi.org/10.1201/b11867-15>
- Bernos, T. A., Travouck, C., Ramasinoro, N., Fraser, D. J., & Mathevon, B. (2021). What can be learned from fishers' perceptions for fishery management planning? Case study insights from Sainte-Marie, Madagascar. *PLoS ONE*, 16(11 November), 1–23. <https://doi.org/10.1371/journal.pone.0259792>
- Bertalanffy, L. (1938). A quantitative theory of organic growth', *Human Biology* (Vol. 10, Issue 2, pp. 181–213). Wayne State University Press. <http://www.jstor.org/stable/41447359>.
- Bethea, D. M., Hale, L., Carlson, J. K., Cortés, E., Manire, C. A., & Gelsleichter, J. (2007). Geographic and ontogenetic variation in the diet and daily ration of the bonnethead shark, *Sphyrna tiburo*, from the eastern Gulf of Mexico. *Marine Biology*, 152(5), 1009–1020. <https://doi.org/10.1007/s00227-007-0728-7>
- Binet, T., Failler, P., Chavance, P. N., & Abidine, M. (2013). First international payment for marine ecosystem services: The case of the Banc d'Arguin National Park, Mauritania. *Global Environmental Change*, 23(6), 1434–1443. <https://doi.org/10.1016/j.gloenvcha.2013.09.015>
- Bird, C. S., Veríssimo, A., Magozzi, S., Abrantes, K. G., Aguilar, A., Al-Reasi, H., & Trueman, C. N. (2018). A global perspective on the trophic geography of sharks. *Nature Ecology & Evolution*, 13–15. <https://doi.org/10.1038/s41559-017-0432-z>

- Bird, E. C. F. (2011). *Coastal geomorphology: An introduction*. John Wiley & Sons.
- Boer, W. F., & Longamane, F. A. (1996). The exploitation of intertidal food resources in inhaca bay, mozambique, by shorebirds and humans. *Biological Conservation*, 78(96), 295–303.
- Boere, G. C., & Stroud, D. A. (2006). The Flyway Concept: What it is and what it isn't. In C. A. Boere, C. A. Galbraith, & D. A. Stroud (Eds.), *Waterbirds around the world* (pp. 40–47). The Stationary Office.
- Boero, F., & Bonsdorff, E. (2007). A conceptual framework for marine biodiversity and ecosystem functioning. *Marine Ecology*, 28, 134–145. <https://doi.org/10.1111/j.1439-0485.2007.00171.x>
- Bom, R. A., Fouw, J., Klaassen, R. H. G., Piersma, T., Lavaleye, M. S. S., Ens, B. J., & Gils, J. A. (2018). Food web consequences of an evolutionary arms race: Molluscs subject to crab predation on intertidal mudflats in Oman are unavailable to shorebirds. *Journal of Biogeography*, 45(2), 342–354. <https://doi.org/10.1111/jbi.13123>
- Booth, H., Squires, D., & Milner-Gulland, E. J. (2019). The neglected complexities of shark fisheries, and priorities for holistic risk-based management. *Ocean and Coastal Management*, 182(July), 104994. <https://doi.org/10.1016/j.ocecoaman.2019.104994>
- Booty, J. M., Underwood, G. J. C., Parris, A., Davies, R. G., & Tolhurst, T. J. (2020). Shorebirds affect ecosystem functioning on an intertidal mudflat. *Frontiers in Marine Science*, 7(August). <https://doi.org/10.3389/fmars.2020.00685>
- Bornatowski, H., Braga, R. R., & Barreto, R. P. (2018). Elasmobranchs consumption in Brazil: Impacts and consequences. In *Advances in marine vertebrate research in latin america* (pp. 251–262). <https://doi.org/10.1007/978-3-319-56985-7>
- Bornatowski, H., Navia, A. F., Braga, R. R., Abilhoa, V., & Corrêa, M. F. M. (2014). Ecological importance of sharks and rays in a structural foodweb analysis in southern Brazil. *ICES J. Mar. Sci*, 71, 1586–1592. <https://doi.org/10.1093/icesjms/fsu025>.
- Bostic, D. L., & Banks, R. C. (1966). A record of stingray predation by the brown pelican. *The Condor*, 68(5), 515–516.
- Botts, R. T., Eppert, A. A., Wiegman, T. J., Rodriguez, A., Blankenship, S. R., Asselin, E. M., & Allen, G. R. (2020). Circadian activity patterns of mammalian predators and prey in Costa Rica. *Journal of Mammalogy*, 101(5), 1313–1331. <https://doi.org/10.1093/jmammal/gyaa103>
- Boulay, S. (2013). Mutations techniques, changements sociaux survenus chez les pêcheurs imrâgen, des années 1970 à nos jours.
- Bouma, T. J., Belzen, J., Balke, T., Zhu, Z., Airoidi, L., Blight, A. J., & Herman, P. M. J. (2014). Identifying knowledge gaps hampering application of intertidal habitats in coastal protection: Opportunities & steps to take. *Coastal Engineering*, 87, 147–157. <https://doi.org/10.1016/j.coastaleng.2013.11.014>
- Boussarie, G., Bakker, J., Wangenstein, O. S., Mariani, S., Bonnin, L., Juhel, J.-B., & Mouillot, D. (2018). Environmental DNA illuminates the dark diversity of sharks. *Science Advances*, 4(5), 9661. <https://doi.org/10.1126/sciadv.aap9661>
- Bowes, R. E., & Thorp, J. H. (2015). Consequences of employing amino acid vs. Bulk-tissue, stable isotope analysis: A laboratory trophic position experiment. *Ecosphere* (Washington, D.C), 6(1). <https://doi.org/10.1890/ES14-00423.1>.
- Braccini, M., Aires-da-Silva, A., & Taylor, I. (2016). Incorporating movement in the modelling of shark and ray population dynamics: Approaches and management implications. *Reviews in Fish Biology and Fisheries*, 26(1), 13–24. <https://doi.org/10.1007/s11160-015-9406-x>
- Braham, C. B., Fréon, P., Laurec, A., Demarcq, H., & Bez, N. (2014). New insights in the spatial dynamics of sardinella stocks off Mauritania (North-West Africa) based on logbook data analysis. *Fisheries Research*, 154, 195–204. <https://doi.org/10.1016/j.fishres.2014.02.020>

- Branco-Nunes, I. S. L., Albuquerque, F. V., Nunes, D. M., Oliveira, P. G. V., & Hazin, F. H. V. (2016). First record of predation between *Dasyatis* species. *Journal of Fish Biology*, 89(4), 2178–2181. <https://doi.org/10.1111/jfb.13091>
- Brandl, S. J., Rasher, D. B., Côté, I. M., Casey, J. M., Darling, E. S., Lefcheck, J. S., & Duffy, J. E. (2019). Coral reef ecosystem functioning: Eight core processes and the role of biodiversity. *Frontiers in Ecology and the Environment*, 17(8), 445–454. <https://doi.org/10.1002/fee.2088>
- Braulik, G., Kasuga, M., & Majubwa, G. (2020). Local ecological knowledge demonstrates shifting baselines and the large-scale decline of sawfishes (*Pristidae*) in Tanzania. *African Journal of Marine Science*, 42(1), 67–79. <https://doi.org/10.2989/1814232X.2020.1728379>
- Bridges, C. R. (1993). Ecophysiology of intertidal fish. In J. C. Rankin & F. B. Jensen (Eds.), *Fish ecophysiology* (pp. 375–400). <https://doi.org/10.1007/978-94-011-2304-415>
- Brinton, C. P., & Curran, M. C. (2017). Tidal and diel movement patterns of the Atlantic stingray (*Dasyatis sabina*) along a stream-order gradient. *Marine and Freshwater Research*.
- Bruno, J. F., Stachowicz, J. J., & Bertness, M. D. (2003). Inclusion of facilitation into ecological theory. *Trends in Ecology and Evolution*, 18(3), 119–125. [https://doi.org/10.1016/S0169-5347\(02\)00045-9](https://doi.org/10.1016/S0169-5347(02)00045-9)
- Buchanan, J. B. (2012). Change in merlin hunting behavior following recovery of peregrine falcon populations suggests mesopredator suppression. *Journal of Raptor Research*, 46(4), 349–356. <https://doi.org/10.3356/JRR-11-75.1>
- Budd, A. M., Cooper, M. K., Port, A. L., Schils, T., Mills, M. S., Deinhart, M. E., & Strugnell, J. M. (2021). First detection of critically endangered scalloped hammerhead sharks (*Sphyrna lewini*). In Guam, micronesia, in five decades using environmental DNA. *Ecological indicators* (Vol. 127). <https://doi.org/10.1016/j.ecolind.2021.107649>
- Buelow, C., & Sheaves, M. (2015). A birds-eye view of biological connectivity in mangrove systems. *Estuarine, Coastal and Shelf Science*, 152, 33–43. <https://doi.org/10.1016/j.ecss.2014.10.014>
- Burgess, M. G., Polasky, S., & Tilman, D. (2013). Predicting overfishing and extinction threats in multispecies fisheries. *Proceedings of the National Academy of Sciences*, 110(40), 15943–15948. <https://doi.org/10.1073/pnas.1314472110>
- Bürkner, P. C. (2018). Advanced bayesian multilevel modeling with the R package brms'. *The R Journal*, 10(July), 395–411.
- Bürkner, P.-C. (2017). brms: An R package for bayesian multilevel models using stan'. *Journal of Statistical Software*, 1(ue 1). <https://www.jstatsoft.org/v080/i01>.
- Buskirk, J., & Yurewicz, K. L. (1998). Effects of predators on prey growth rate: Relative contributions of thinning and reduced activity. *Oikos* (Copenhagen, Denmark), 82(1), 20. <https://doi.org/10.2307/3546913>
- Byers J. E., & Grabowski J. H. (2014). Soft sediment communities. In Bertness M. D., Bruno J. F., Silliman B. R., & Stachowicz J. J. (Eds.), *Marine Community Ecology* (pp. 227–249). <https://doi.org/10.1007/978-94-017-8801-4151>

C

- Cadée, G. C. (2001). Sediment Dynamics by Bioturbating Organisms. In *Ecological Comparisons of Sedimentary Shores* (Vol. 151, pp. 127–148). <https://doi.org/10.1007/978-3-642-56557-17>
- Campana, S. E. (2014). Age determination of elasmobranchs with special reference to mediterranean species., GFCM. *Studies and Reviews*. <http://www.fao.org/documents/card/en/c/48294ee7-cd21-4388-a51e-6f381035f407/>.

- Campos, B. R., Fish, M. A., Jones, G., Riley, R. W., Allen, P. J., Klimley, P. A., & Kelly, J. T. (2009). Movements of brown smoothhounds, *Mustelus henlei*, in Tomales Bay, California. *Environmental Biology of Fishes*, 85(1), 3–13. <https://doi.org/10.1007/s10641-009-9462-y>
- Campredon, P., & Catry, P. (2016). Bijagós Archipelago (Guinea-Bissau). In *The Wetland Book* (pp. 1–8). Springer Netherlands.
- Cardinale, B. J., Duffy, J. E., Gonzalez, A., Hooper, D. U., Perrings, C., Venail, P., & Naeem, S. (2012). Biodiversity loss and its impact on humanity. *Nature*, 489(7415), 326–326. <https://doi.org/10.1038/nature11373>
- Cardinale, M., Chacate, O., Casini, M., Chaúca, I., & Helge Vølstad, J. (2014). CPUE trends of *Hilsa kelee* and *Thryssa vitrirostris* exploited by the artisanal finfish fisheries in Mozambique derived from an on-shore sampling of catches by trip. *Sci. Mar*, 78, 55–64. <https://doi.org/10.3989/scimar.03826.06c>
- Carlier, A., Chauvaud, L., Geest, M., Le Loc'h, F., Le Duff, M., Vernet, M., & Clavier, J. (2015). Trophic connectivity between offshore upwelling and the inshore food web of Banc d'Arguin (Mauritania): New insights from isotopic analysis. *Estuarine, Coastal and Shelf Science*, 165(August), 149–158. <https://doi.org/10.1016/j.ecss.2015.05.001>
- Carlisle, A. B., & Starr, R. M. (2009). Habitat use, residency, and seasonal distribution of female leopard sharks *Triakis semifasciata* in Elkhorn Slough, California. *Marine Ecology Progress Series*, 380(June 2014), 213–228. <https://doi.org/10.3354/meps07907>
- Carlisle, A. B., & Starr, R. M. (2010). Tidal movements of female leopard sharks (*Triakis semifasciata*) in Elkhorn Slough, California. *Environmental Biology of Fishes*, 89(1), 31–45. <https://doi.org/10.1007/s10641-010-9667-0>
- Carlson, J., Charvet, P., Blanco-Parra, M. P., Briones Bell-Iloch, A., Cardenosa, D., Derrick, D., Espinoza, E., Morales-Saldaña, J. M., Naranjo-Elizondo, B., Pacoureaux, N., Schneider, E. V. C., Simpson, N. J., Pollom, R., & Dulvy, N. K. (2020). *Hypanus americanus*. In *The IUCN Red List of Threatened Species 2020* (p. 181244884 104123787). <https://doi.org/10.2305/IUCN.UK.2020-3.RLTS.T181244884A104123787.en>
- Carlton, J. T., & Hodder, J. (2003). Maritime mammals: Terrestrial mammals as consumers in marine intertidal communities. *Marine Ecology Progress Series*, 256, 271–286.
- Casey, J. M., Baird, A. H., Brandl, S. J., Hoogenboom, M. O., Rizzari, J. R., Frisch, A. J., & Connolly, S. R. (2017). A test of trophic cascade theory: Fish and benthic assemblages across a predator density gradient on coral reefs. *Oecologia*, 183(1), 161–175. <https://doi.org/10.1007/s00442-016-3753-8>
- Castellanos-Galindo, G. A., Herrón, P., Navia, A. F., & Booth, H. (2021). Shark conservation and blanket bans in the eastern Pacific Ocean. *Conservation Science and Practice*, 3(7), 2–5. <https://doi.org/10.1111/csp2.428>
- Castilla, J. C. (1998). Coastal marine communities: Trends and perspectives from human-exclusion experiments. *Trends in Ecology and Evolution*, 14(7), 280–283. [https://doi.org/10.1016/S0169-5347\(99\)01602-X](https://doi.org/10.1016/S0169-5347(99)01602-X)
- Castillo-Géniz, J. L., Márquez-Farías, J. F., Cruz, M. C., Cortés, E., & Cid Del Prado, A. (1998). The Mexican artisanal shark fishery in the Gulf of Mexico: Towards a regulated fishery. *Marine and Freshwater Research*, 49(7), 611–620. <https://doi.org/10.1071/MF97120>
- Castorani, M. C. N., & Hovel, K. A. (2016). Native predator chemical cues induce anti-predation behaviors in an invasive marine bivalve. *Biological Invasions*, 18(1), 169–181. <https://doi.org/10.1007/s10530-015-1000-6>
- Castro, J. I. (1996). Biology of the blacktip shark, *Carcharhinus limbatus*. *Bulletin of Marine Science*, 59(3), 508–522.

- Catry, P., Barbosa, C., Indjai, B., Almeida, A., Godley, B. J., & Vié, J. C. (2002). First census of the green turtle at Poilão, Bijagós Archipelago, Guinea-Bissau: The most important nesting colony on the Atlantic coast of Africa. *Oryx: The Journal of the Fauna Preservation Society*, 36(4), 400–403. <https://doi.org/10.1017/S0030605302000765>
- Catry, T., Lourenço, P. M., Lopes, R. J., Carneiro, C., Alves, J. A., Costa, J., & Granadeiro, J. P. (2016). Structure and functioning of intertidal food webs along an avian flyway: A comparative approach using stable isotopes. *Functional Ecology*, 30(3), 468–478. <https://doi.org/10.1111/1365-2435.12506>
- Caut, S., Angulo, E., & Courchamp, F. (2009). Variation in discrimination factors ($\Delta^{15}\text{N}$ and $\Delta^{13}\text{C}$): The effect of diet isotopic values and applications for diet reconstruction. *Journal of Applied Ecology*, 46(2), 443–453. <https://doi.org/10.1111/j.1365-2664.2009.01620.x>
- Cerutti-Pereyra, F., Thums, M., Austin, C. M., Bradshaw, C. J. A., Stevens, J. D., Babcock, R. C., & Meekan, M. G. (2014). Restricted movements of juvenile rays in the lagoon of Ningaloo Reef, Western Australia—Evidence for the existence of a nursery. *Environmental Biology of Fishes*, 97(4), 371–383. <https://doi.org/10.1007/s10641-013-0158-y>
- Chapman, D. D., & Gruber, S. H. (2002). A further observation of the prey-handling behavior of the great hammerhead shark, *Sphyrna mokarran*: Predation upon the spotted eagle ray, *Aetobatus Narinari*. *Bulletin of Marine Science*, 70(3), 947–952.
- Chew, S. F., Wilson, J. M., Ip, Y. K., & Randall, D. J. (2006). Nitrogen excretion and defense against ammonia toxicity. In A. L. Val, V. M. F. Almeida-Val, & D. J. Randall (Eds.), *The physiology of tropical fishes* (pp. 307–396). Academic Press.
- Chin, A., Heupel, M. R., Simpfendorfer, C. A., & Tobin, A. J. (2016). Population organisation in reef sharks: New variations in coastal habitat use by mobile marine predators. *Marine Ecology Progress Series*, 544, 197–211. <https://doi.org/10.3354/meps11545>
- Chuenpagdee, R., Liguori, L., Palomares, M. L. D., & Pauly, D. (2006). Bottom-up, global estimates of small-scale marine fisheries catches. *Fisheries Centre Research Reports*, 14(8), 105.
- Cisneros-Montemayor, A. M., Barnes-Mauthe, M., Al-Abdulrazzak, D., Navarro-Holm, E., & Sumaila, U. R. (2013). Global economic value of shark ecotourism: Implications for conservation. *Oryx: The Journal of the Fauna Preservation Society*, 47(3), 381–388. <https://doi.org/10.1017/S0030605312001718>
- Clarke, K. R., & Green, R. H. (1988). Statistical design and analysis for a “biological effects” study. *Marine Ecology - Progress Series*, 46, 213–226.
- Clarke, S., Milner-Gulland, E. J., & Trond, B. (2007). Social, economic, and regulatory drivers of the shark fin trade. *Marine Resource Economics*, 22(3), 305–327. <https://doi.org/10.1086/mre.22.3.42629561>
- Clements, O. N., Leurs, G., Witbaard, R., Pen, I., Verkuil, Y. I., & Govers, L. L. (2022). Growth, maturity, and diet of the pearl whipray (*Fontitrygon margaritella*) from the Bijagós Archipelago, Guinea-Bissau. *PeerJ*, 10, 12894. <https://doi.org/10.7717/peerj.12894>
- Cliff, G. (1995). Sharks caught in the protective gill nets off KwaZulu-Natal, South Africa. 8. The Great hammerhead shark *Sphyrna mokarran*. *South African Journal of Marine Science*, 15(1), 105–114. <https://doi.org/10.2989/025776195784156331>
- Cliff, G., & Dudley, S. F. (1991). Sharks caught in the protective gill nets off Natal, South Africa. 4. The bull shark *Carcharhinus leucas*. *South African Journal of Marine Science*, 10(1), 253–270. <https://doi.org/10.2989/02577619109504636>
- Cliff, G., Dudley, S. F. J., & Davis, B. (1989). Sharks caught in the protective gill nets off Natal, South Africa. 2. The great white shark *Carcharodon carcharias*. *South African Journal of Marine Science*, 8(1), 131–144. <https://doi.org/10.2989/02577618909504556>

- Coelho, A. P., Henriques, M., Rocha, A. D., Paulino, J., Schaars, L. K., Ramos, C., & Alves, J. A. (2022). Spatial and seasonal variation in macrozoobenthic density, biomass and community composition in a major tropical intertidal area, the Bijagós Archipelago, West Africa. *PLoS ONE*, 17(11 November), 1–27. <https://doi.org/10.1371/journal.pone.0277861>
- Cohen, J. E., Pimm, S. L., Yodzis, P., & Saldaña, J. (1993). Body sizes of animal predators and animal prey in food webs. *The Journal of Animal Ecology*, 62, 67–78.
- Coiraton, C., Amezcua, F., & Ketchum, J. T. (2020). New insights into the migration patterns of the scalloped hammerhead shark *Sphyrna lewini* based on vertebral microchemistry. *Marine Biology*, 167(5), 1–18. <https://doi.org/10.1007/s00227-020-3668-0>
- Coll, M., Navarro, J., & Palomera, I. (2013). Ecological role, fishing impact, and management options for the recovery of a Mediterranean endemic skate by means of food web models'. *Biological Conservation*. Elsevier Ltd, 157, 108–120. <https://doi.org/10.1016/j.biocon.2012.06.029>.
- Collins, A. B., Heupel, M. R., Hueter, R. E., & Motta, P. J. (2007). Hard prey specialists or opportunistic generalists? An examination of the diet of the cownose ray, *Rhinoptera bonasus*. *Marine and Freshwater Research*, 58(1), 135–144. <https://doi.org/10.1071/MF05227>
- Colloca, F., Carrozzi, V., Simonetti, A., & Lorenzo, M. (2020). Using local ecological knowledge of fishers to reconstruct abundance trends of elasmobranch populations in the Strait of Sicily. *Frontiers in Marine Science*, 7(June), 1–8. <https://doi.org/10.3389/fmars.2020.00508>
- Compagno, L. J. V., & Roberts, T. R. (1984). Marine and freshwater stingrays (Dasyatidae) of West Africa, with description of a new species. *Proceedings of the California Academy of Sciences*, 43(18), 283–300.
- Compton, T. J., Holthuijsen, S., Koolhaas, A., Dekinga, A., Horn, J., Smith, J., Galama, Y., Brugge, M., Wal, D., Meer, J., Veer, H. W., & Piersma, T. (2013). Distinctly variable mudscapes: Distribution gradients of intertidal macrofauna across the Dutch Wadden Sea. *Journal of Sea Research*, 82, 103–116.
- Condit, R., & Le Boeuf, B. J. (1984). Feeding Habits and Feeding Grounds of the Northern Elephant Seal. *Journal of Mammalogy*, 65(2), 281–290. <https://doi.org/10.2307/1381167>
- Conrath, C. L., & Musick, J. A. (2010). Residency, space use and movement patterns of juvenile sandbar sharks (*Carcharhinus plumbeus*) within a Virginia summer nursery area. *Marine and Freshwater Research*, 61(2), 223–235. <https://doi.org/10.1071/MF09078>
- Convertino, M., Donoghue, J. F., Chu-Agor, M. L., Kiker, G. A., Muñoz-Carpena, R., Fischer, R. A., & Linkov, I. (2011). Anthropogenic renourishment feedback on shorebirds: A multispecies Bayesian perspective. *Ecological Engineering*, 37(8), 1184–1194. <https://doi.org/10.1016/j.ecoleng.2011.02.019>
- Conway, J. N., & Mcfee, W. E. (2017). Ingestion of stingrays (*Dasyatis* spp.) by a common bottlenose dolphin (*Tursiops truncatus*). *Aquatic Mammals*, 1–10.
- Corgos, A., & Rosende-Pereiro, A. (2022). Nursery habitat use patterns of the scalloped hammerhead shark, *Sphyrna lewini*, in coastal areas of the central Mexican Pacific. *Journal of Fish Biology*, 100(1), 117–133. <https://doi.org/10.1111/jfb.14925>
- Correia, E., Granadeiro, J. P., Regalla, A., & Catry, P. (2021). Coastal waters of a marine protected area of the Bijagós Archipelago, West Africa, shelter juvenile fishes of economic and ecological importance. *Regional Studies in Marine Science*, 46, 101892. <https://doi.org/10.1016/j.rsma.2021.101892>
- Correia, E., Granadeiro, J. P., Santos, B., Regalla, A., Mata, V. A., & Catry, T. (2023). Trophic ecology of a migratory shorebird community at a globally important non-breeding site: Combining DNA metabarcoding and conventional techniques. *Marine Ecology Progress Series*, 705, 127–144.

- Cortés, E., & Gruber, S. H. (1990). Diet, feeding habits and estimates of daily ration of young lemon sharks, *Negaprion brevirostris*. *Copeia*, 1990(1), 204. <https://doi.org/10.2307/1445836>
- Cowley, P. D. (1997). Age and growth of the blue stingray *Dasyatis chrysonota chrysonota* from the South-Eastern Cape coast of South Africa'. *South African Journal of Marine Science*, 7615(18), 31–38. <https://doi.org/10.2989/025776197784161054>.
- Cox, D. L., & Koob, T. J. (1993). Predation on elasmobranch eggs. *Environmental Biology of Fishes*, 38, 117–125. <https://doi.org/10.1007/978-94-017-3450-910>
- Crain, C. M., Halpern, B. S., Beck, M. W., & Kappel, C. V. (2009). Understanding and managing human threats to the coastal marine environment. In *Annals of the new york academy of sciences* (pp. 39–62). <https://doi.org/10.1111/j.1749-6632.2009.04496.x>
- Cross, H. (2015). Elasmobranch capture by commercial small-scale fisheries in the Bijagós Archipelago, Guinea-Bissau. *Fisheries Research*, 168, 105–108. <https://doi.org/10.1016/j.fishres.2015.03.018>
- Cross, H. C. (2014). The importance of small-scale fishing to rural coastal livelihoods: A comparative case-study in the Bijagós Archipelago Guinea-Bissau [University College London]. <https://doi.org/10.1007/s13398-014-0173-7.2>
- Curras, M. R., Donadio, E., Middleton, A. D., & Pauli, J. N. (2022). Carnivore niche partitioning in a human landscape. *American Naturalist*, 199(4). <https://doi.org/10.1086/718472>
- Curtis, T. H., & Macesic, L. J. (2011). Observations of breaching behavior in juvenile bull sharks, *Carcharhinus leucas*. *Florida Scientist*, 74(4), 253–257.
- Cushing, D. H. (1971). Upwelling and the production of fish. *Advances in Marine Biology*, 9, 255–334.

D

- Dabruzzi, T. F., Bennett, W. A., Rummer, J. L., & Fanguie, N. A. (2013). Juvenile Ribbontail Stingray, *Taeniura lymma* (Forsskål, 1775) (Chondrichthyes, Dasyatidae), demonstrate a unique suite of physiological adaptations to survive hyperthermic nursery conditions. *Hydrobiologia*, 701(1), 37–49. <https://doi.org/10.1007/s10750-012-1249-z>
- Dale, J. J., & Holland, K. N. (2012). Age, growth and maturity of the brown stingray (*Dasyatis lata*) around Oahu, Hawai'i. *Marine and Freshwater Research*, 63(6), 475–484. <https://doi.org/10.1071/MF11231>.
- Daly, R., Froneman, P. W., & Smale, M. J. (2013). Comparative feeding ecology of bull sharks (*Carcharhinus leucas*) in the coastal waters of the Southwest Indian Ocean inferred from stable isotope analysis. *PLoS ONE*, 8(10), 1–11. <https://doi.org/10.1371/journal.pone.0078229>
- Daly, R., Smale, M. J., Singh, S., Anders, D., Shivji, M., Clare, C. A., & Barnett, A. (2018). Refuges and risks: Evaluating the benefits of an expanded MPA network for mobile apex predators. *Diversity and Distributions*, 24(9), 1217–1230. <https://doi.org/10.1111/ddi.12758>
- Daly-Engel, T. S., Seraphin, K. D., Holland, K. N., Coffey, J. P., Nance, H. a, Toonen, R. J., & Bowen, B. W. (2012). Global phylogeography with mixed-marker analysis reveals male-mediated dispersal in the endangered scalloped hammerhead shark (*Sphyrna lewini*). *PLoS ONE*, 7(1). <https://doi.org/10.1371/journal.pone.0029986>
- D'Andrea, A. F., Aller, R. C., & Lopez, G. R. (2002). Organic matter flux and reactivity on a South Carolina sandflat: The impacts of porewater advection and macrobiological structures. *Limnology and Oceanography*, 47, 1056–1070.

- D'Andrea, A. F., Lopez, G. R., & Aller, R. C. (2004). Rapid physical and biological particle mixing on an intertidal sandflat. *Journal of Marine Research*, 62(1), 67–92. <https://doi.org/10.1357/00222400460744627>
- Darimont, C. T., Cooke, R., Bourbonnais, M. L., Bryan, H. M., Carlson, S. M., Estes, J. A., & Maclean, J. L. (2023). Humanity's diverse predatory niche and its ecological consequences. *Communications Biology*. <https://doi.org/10.1038/s42003-023-04940-w>
- Darimont, C. T., Fox, C. H., Bryan, H. M., & Reimchen, T. E. (2015). The unique ecology of human predators. *Science (New York, N.Y.)*, 349(6250), 858–860. <https://doi.org/10.1126/science.aac4249>
- Davy, L. E., Simpfendorfer, C. A., & Heupel, M. R. (2015). Movement patterns and habitat use of juvenile mangrove whiphrays (*Himantura granulata*) (pp. 481–492). *Marine and Freshwater Research*.
- Day, J., Clark, J. A., Williamson, J. E., Brown, C., & Gillings, M. (2019). Population genetic analyses reveal female reproductive philopatry in the oviparous Port Jackson shark. *Marine and Freshwater Research*, 70(7), 986–994. <https://doi.org/10.1071/MF18255>
- Deagle, B. E., Kirkwood, R., & Jarman, S. N. (2009). Analysis of Australian fur seal diet by pyrosequencing prey DNA in faeces. *Molecular Ecology*, 18(9), 2022–2038. <https://doi.org/10.1111/j.1365-294X.2009.04158.x>
- Dean, M. N., Bizzarro, J. J., Clark, B., Underwood, C. J., & Johanson, Z. (2017). Large batoid fishes frequently consume stingrays despite skeletal damage. *Royal Society Open Science*, 4(9). <https://doi.org/10.1098/rsos.170674>
- Debaere, S. F., Weideli, O. C., Bouyoucos, I. A., Eustache, K. B., Trujillo, J. E., Boeck, G., & Stacy, N. (2023). Quantifying changes in umbilicus size to estimate the relative age of neonatal blacktip reef sharks (*Carcharhinus melanopterus*). *Conservation Physiology*, 11(1), 1–11. <https://doi.org/10.1093/conphys/coad028>
- Deegan, L. A., Johnson, D. S., Warren, R. S., Peterson, B. J., Fleeger, J. W., Fagherazzi, S., & Wollheim, W. M. (2012). Coastal eutrophication as a driver of salt marsh loss. *Nature*, 490(7420), 388–392. <https://doi.org/10.1038/nature11533>
- Den Hout, P. J., Spaans, B., & Piersma, T. (2008). Differential mortality of wintering shorebirds on the Banc d'Arguin, Mauritania, due to predation by large falcons. *Ibis*, 150(SUPPL.1), 219–230. <https://doi.org/10.1111/j.1474-919X.2008.00785.x>
- Dent, F., & Clarke, S. (2015). State of the global market for shark products. *FAO Technical Paper*, 196. <http://www.fao.org/docrep/field/003/ab825f/AB825F00.htm#TOC>
- Deppe, F. (1999). *Intertidal Mudflats Worldwide*. Common Wadden Sea Secretariat.
- Derrick, B., Burns, K., Zhu, A., Andreoli, V., Zeller, D., & Pauly, D. (2023). Small-scale fisheries catch and fishing effort in the Socotra Archipelago (Yemen) between 1950 and 2019. *Frontiers in Marine Science*, 10(June), 1–15. <https://doi.org/10.3389/fmars.2023.1201661>
- Derrick, D. H., Cheok, J., & Dulvy, N. K. (2020). Spatially congruent sites of importance for global shark and ray biodiversity. *PLoS ONE*, 15(7 July), 1–19. <https://doi.org/10.1371/journal.pone.0235559>
- Dhanjal-Adams, K. L., Hanson, J. O., Murray, N. J., Phinn, S. R., Wingate, V. R., Mustin, K., Lee, J. R., Allan, J. R., Cappadonna, J. L., Studds, C. E., Clemens, R., Roelfsema, C. M., & Fuller, R. A. (2016). The distribution and protection of intertidal habitats in Australia. *Emu - Austral Ornithology*, 116(2), 208–214. <https://doi.org/10.1071/MU15046>
- Dicken, M. L., Hussey, N. E., Christiansen, H. M., Smale, M. J., Nkabi, N., Cliff, G., & Wintner, S. P. (2017). Diet and trophic ecology of the tiger shark (*Galeocerdo cuvier*) from South African waters. *PLoS ONE*, 12(6), 1–25. <https://doi.org/10.1371/journal.pone.0177897>

- Diemer, K. M., Mann, B. Q., & Hussey, N. E. (2011). Distribution and movement of scalloped hammerhead *Sphyrna lewini* and smooth hammerhead *Sphyrna zygaena* sharks along the east coast of Southern Africa. *African Journal of Marine Science*, 33(2), 229–238. <https://doi.org/10.2989/1814232X.2011.600291>
- Dinerstein, E., Vynne, C., Sala, E., Joshi, A. R., Fernando, S., Lovejoy, T. E., & Wikramanayake, E. (2019). A global deal for nature: Guiding principles, milestones, and targets. *Science Advances*, 5(4), 1–18. <https://doi.org/10.1126/sciadv.aaw2869>
- Diop, M., & Dossa, J. (2011). 30 Years of shark fishing. *IUCN Shark Spec. Gr*, 51. <http://www.iucnssg.org/uploads/5/4/1/2/54120303/30yearseng.pdf>.
- Diouf, P. S., Deme-Gningue, I., & Albaret, J. J. (1994). L'Archipel des Bijagós environment aquatique et peuplement de poissons. *Ministere Des Peches de Guinee Bissau*.
- Ditlevsen, P. D., & Ditlevsen, S. (2023). Warning of a forthcoming collapse of the Atlantic meridional overturning circulation. *Nature Communications*, 14(4254), 1–12. <https://doi.org/10.1038/s41467-023-39810-w>
- Donald, P. F., Fishpool, L. D. C., Ajagbe, A., Bennun, L. A., Bunting, G., Burfield, I. J., & Wege, D. C. (2019). Important Bird and Biodiversity Areas (IBAs): The development and characteristics of a global inventory of key sites for biodiversity. *Bird Conservation International*, 29(2), 177–198. <https://doi.org/10.1017/S0959270918000102>
- Dubay, K., Tokuoka, S., & Gereffi, G. (2010). A Value Chain Analysis of the Sinaloa. *Mexico Shrimp Fishery*, 919.
- Ducrocq, M., Lemine Ould Sidi, M., & Ould Yarba, L. (2004). Comment le Parc national du Banc d'Arguin est devenu le plus grand sanctuaire d'Afrique pour les requins. *FIBA*.
- Dudley, S. F., & Cliff, G. (1993). Sharks caught in the protective gill nets off Natal, South Africa. 7. The blacktip shark *Carcharhinus limbatus*. *South African Journal of Marine Science*, 13(1), 237–254. <https://doi.org/10.2989/025776193784287356>
- Dulvy, N. K., Fowler, S. L., Musick, J. A., Cavanagh, R. D., Kyne, P. M., Harrison, L. R., Carlson, J. K., Davidson, L. N., Fordham, S. V., Francis, M. P., Pollock, C. M., Simpfendorfer, C. A., Burgess, G. H., Carpenter, K. E., Compagno, L. J., Ebert, D. A., Gibson, C., Heupel, M. R., Livingstone, S., SR, ... W.T. (2014). Extinction risk and conservation of the world's sharks and rays. *Elife*, 3:e00590.
- Dulvy, N. K., Metcalfe, J. D., Glanville, J., Pawson, M. G., & Reynolds, J. D. (2000). Fishery stability, local extinctions, and shifts in community structure in skates. *Conservation Biology*, 14(1), 283–293. <https://doi.org/10.1046/j.1523-1739.2000.98540.x>
- Dulvy, N. K., Pacoureau, N., Rigby, C. L., Pollom, R. A., Jabado, R. W., Ebert, D. A., Finucci, B., Pollock, C. M., Cheok, J., Derrick, D. H., Herman, K. B., Sherman, C. S., VanderWright, W. J., Lawson, J. M., Walls, R. H. L., Carlson, J. K., Charvet, P., Bineesh, K. K., Fernando, D., ... Simpfendorfer, C. A. (2021). Overfishing drives over one-third of all sharks and rays toward a global extinction crisis. *Current Biology* : CB, 31, 4773–4787.
- Dulvy, N. K., Sadovy, Y., & Reynolds, J. D. (2003). Extinction vulnerability in marine populations. *Fish and Fisheries*, 4(1), 25–64. <https://doi.org/10.1046/j.1467-2979.2003.00105.x>
- Dulvy, N. K., Simpfendorfer, C. A., Davidson, L. N. K., Fordham, S. V., Bräutigam, A., Sant, G., & Welch, D. J. (2017). Challenges and priorities in shark and ray conservation. *Current Biology*, 27(11), 565–572. <https://doi.org/10.1016/j.cub.2017.04.038>
- Dunn, N., Savolainen, V., Weber, S., Andrzejczek, S., Carbone, C., & Curnick, D. (2022). Elasmobranch diversity across a remote coral reef atoll revealed through environmental DNA metabarcoding. *Zoological Journal of the Linnean Society*, 1–15. <https://doi.org/10.1093/zoolinnean/zlac014>

E

- Early-Capistrán, M. M., Solana-Arellano, E., Alberto Abreu-Grobois, F., Narchi, N. E., Garibay-Melo, G., Seminoff, J. A., & Saenz-Arroyo, A. (2020). Quantifying local ecological knowledge to model historical abundance of long-lived, heavily-exploited fauna. *PeerJ*, 8, 1–34. <https://doi.org/10.7717/peerj.9494>
- Ebert, D. A., Dando, M., & Fowler, S. (2021). *Sharks of the world: A complete guide*. Princeton University Press.
- Eckert, S. A., Bagley, D., Kubis, S., Ehrhart, L., Johnson, C., Stewart, K., & DeFreese, D. (2006). Internesting and postnesting movements and foraging habitats of leatherback sea turtles (*Dermochelys coriacea*) nesting in Florida. *Chelonian Conservation and Biology*, 5(2), 239–248. [https://doi.org/10.2744/1071-8443\(2006\)5](https://doi.org/10.2744/1071-8443(2006)5)
- Edgar, R. C. (2010). Search and clustering orders of magnitude faster than BLAST. *Bioinformatics (Oxford, England)*, 26(19), 2460–2461. <https://doi.org/10.1093/bioinformatics/btq461>
- Eichmiller, J. J., Best, S. E., & Sorensen, P. W. (2016). Effects of temperature and trophic state on degradation of environmental DNA in lake water. *Environmental Science and Technology*, 50(4), 1859–1867. <https://doi.org/10.1021/acs.est.5b05672>
- Ekpo, I., & Essien-Ibok, M. (2019). Development, prospects and challenges of artisanal fisheries in Akwa Ibom State, Nigeria. *International Journal of Environmental Science, Management and Engineering Research*, 2(3), 69–86.
- El-Hacen, E. H. M., Bouma, T. J., Govers, L. L., Piersma, T., & Olff, H. (2019). Seagrass sensitivity to collapse along a hydrodynamic gradient: Evidence from a pristine subtropical intertidal ecosystem. *Ecosystems (New York, N.Y.)*, 22(5), 1007–1023. <https://doi.org/10.1007/s10021-018-0319-0>
- El-Hacen, E. H. M., Bouma, T. J., Oomen, P., Piersma, T., & Olff, H. (2019). Large-scale ecosystem engineering by flamingos and fiddler crabs on West African intertidal flats promote joint food availability. *Oikos (Copenhagen, Denmark)*, 128, 753–764.
- El-Hacen, E. H. M., Sidi Cheikh, M. A., Bouma, T. J., Olff, H., & Piersma, T. (2020). Long-term changes in seagrass and benthos at Banc d'Arguin, Mauritania, the premier intertidal system along the East Atlantic Flyway. *Global Ecology and Conservation*, 24, 01364. <https://doi.org/10.1016/j.gecco.2020.e01364>
- El-Hacen, E., Lemrabott, S., Meijer, K., Piersma, T., Govers, L., Gils, J., & Olff, H. (2023). Growth and population structure of bloody cockles *Senilia senilis* at Banc d'Arguin and Bijagós with different environmental conditions and harvesting regimes. *Marine Ecology Progress Series*, 710, 71–83. <https://doi.org/10.3354/meps14291>
- Ellis, W. L., & Bell, S. S. (2008). Tidal influence on a fringing mangrove intertidal fish community as observed by in situ video recording: Implications for studies of tidally migrating nekton. *Marine Ecology Progress Series*, 370(November 2015), 207–219. <https://doi.org/10.3354/meps07567>
- Elston, C., Cowley, P. D., Brandis, R. G., & Lea, J. (2021). Residency and habitat use patterns by sympatric stingrays at a remote atoll in the Western Indian Ocean. *Marine Ecology Progress Series*, 662, 97–114. <https://doi.org/10.3354/meps13632>
- Elston, C., Cowley, P. D., Brandis, R. G., & Lea, J. (2022). Stingray habitat use is dynamically influenced by temperature and tides. *Frontiers in Marine Science*, 8(January), 1–13. <https://doi.org/10.3389/fmars.2021.754404>
- Ens, B. J., Piersma, T., & Drent, R. H. (1994). The dependence of waders and waterfowl migrating along the East Atlantic Flyway on their coastal food supplies. *Ophelia*, 6, 127–151.

- Estes, J. A., Heithaus, M., McCauley, D. J., Rasher, D. B., & Worm, B. (2016). Megafaunal impacts on structure and function of ocean ecosystems. *Annual Review of Environment and Resources*, 41. <https://doi.org/10.1146/annurev-environ-110615-085622>
- Estes, J. A., Terborgh, J., Brashares, J. S., Power, M. E., Berger, J., Bond, W. J., & Wardle, D. A. (2011). Trophic downgrading of planet earth. *Science (New York, N.Y.)*, 333(6040), 301–306. <https://doi.org/10.1126/science.1205106>
- Estes, J. A., Tinker, M. T., Williams, T. M., & Doak, D. F. (1998). Killer whale predation on sea otters linking oceanic and nearshore ecosystems. *Science (New York, N.Y.)*, 282(5388), 473–476. <https://doi.org/10.1126/science.282.5388.473>
- Estupiñán-Montaño, C., Estupiñán-Ortiz, J. F., Cedeño-Figueroa, L. G., Galván-Magaña, F., & Polo-Silva, C. J. (2017). Diet of the bull shark, *Carcharhinus leucas*, and the tiger shark, *Galeocerdo cuvier*, in the eastern Pacific Ocean. *Turkish Journal of Zoology*, 41(6), 1111–1117. <https://doi.org/10.3906/zoo-1610-31>
- Estupiñán-Montaño, C., Pacheco-Triviño, F., Cedeño-Figueroa, L. G., Galván-Magaña, F., & Estupiñán-Ortiz, J. F. (2018). Diet of three shark species in the Ecuadorian Pacific, *Carcharhinus falciformis*, *Carcharhinus limbatus* and *Nasolamia velox*. *Journal of the Marine Biological Association of the United Kingdom*, 98(4), 927–935. <https://doi.org/10.1017/S002531541600179X>
- F**
- FAO. (n.d.). Database of measures on conservation and management of sharks. FAO. <https://www.fao.org/ipoa-sharks/database-of-measures/en/>
- FAO. (1999). International Plan of Action for reducing incidental catch of seabirds in longline fisheries. International Plan of Action for the conservation and management of sharks. International Plan of Action for the Management of Fishing Capacity. Rome. <https://doi.org/10.1163/9789004502789033>
- FAO. (2019). Report of the FAO working group on the assessment of small pelagic fish off Northwest Africa.
- F.A.O. (2022). The state of the world fisheries and aquaculture 2022. Towards Blue Transformation. <https://doi.org/10.4060/cc0463en>
- Faria, F. A., Albertoni, E. F., & Bugoni, L. (2018). Trophic niches and feeding relationships of shorebirds in southern Brazil. *Aquatic Ecology*, 52(4), 281–296. <https://doi.org/10.1007/s10452-018-9663-6>
- Farias, I. (2006). Diet comparison of four ray species (*Raja clavata*, *Raja brachyura*, *Raja montagui* and *Leucoraja naevus*) caught along the Portuguese continental shelf. *Aquatic Living Resources*, 19(2), 105–114. <https://doi.org/10.1051/alr:2006010>.
- Feldheim, K. A., Gruber, S. H., Dibattista, J. D., Babcock, E. A., Kessel, S. T., Hendry, A. P., & Chapman, D. D. (2014). Two decades of genetic profiling yields first evidence of natal philopatry and long-term fidelity to parturition sites in sharks. *Molecular Ecology*, 23(1), 110–117. <https://doi.org/10.1111/mec.12583>
- Fernandes, P. G., Ralph, G. M., Nieto, A., Criado, M. G., Vasilakopoulos, P., & Maravelias, C. D. (2017). Coherent assessments of Europe's marine fishes show regional divergence and megafauna loss. *Nature Ecology & Evolution*, 1, 1–16. <https://doi.org/10.1038/s41559-017-0200>.
- Fernández, L., Salmerón, F., & Ramos, A. (2005). Change in elasmobranchs and other incidental species in the Spanish deepwater black hake trawl fishery off Mauritania (1992–2001). *J. Northwest Atl. Fish. Sci*, 35, 325–331. <https://doi.org/10.2960/J.v35.m534>.

- Fernandez-Carvalho, J., Coelho, R., Santos, M. N., & Amorim, S. (2015). Effects of hook and bait in a tropical northeast Atlantic pelagic longline fishery: Part II-Target, bycatch and discard fishes. *Fisheries Research*, 164(3), 312–321. <https://doi.org/10.1016/j.fishres.2014.11.009>
- Fernández-Ordóñez, J. C., Lasso-alcá, O. M., & Ron, E. J. (2016). First evidence of an Osprey (*Pandion haliaetus*) preying on elasmobranches. *Revista Venezolana de Ornitología*, 6, 65–67.
- Fernando, D., & Stewart, J. D. (2021). High bycatch rates of manta and devil rays in the “small-scale” artisanal fisheries of Sri Lanka. *PeerJ*, 9, 1–35. <https://doi.org/10.7717/peerj.11994>
- Ferreira, L. C., Thums, M., Heithaus, M. R., Barnett, A., Abrantes, K. G., Holmes, B. J., & Meekan, M. G. (2017). The trophic role of a large marine predator, the tiger shark *Galeocerdo cuvier*. *Scientific Reports*, 7(1), 7641. <https://doi.org/10.1038/s41598-017-07751-2>
- Ferretti, F., Worm, B., Britten, G. L., Heithaus, M. R., & Lotze, H. K. (2010). Patterns and ecosystem consequences of shark declines in the ocean. *Ecology Letters*, 13(8), 1055–1071. <https://doi.org/10.1111/j.1461-0248.2010.01489.x>
- Fezzi, C., Ford, D. J., & Oleson, K. L. (2023). The economic value of coral reefs: Climate change impacts and spatial targeting of restoration measures. *Ecological Economics*, 203(March 2022), 107628. <https://doi.org/10.1016/j.ecolecon.2022.107628>
- Finke, D. L., & Denno, R. F. (2004). Predator diversity dampens trophic cascades. *Nature*, 429(6990), 407–410. <https://doi.org/10.1038/nature02554>
- Fisher, R. A., Call, G. C., & Grubbs, R. D. (2011). Cownose ray (*Rhinoptera bonasus*) predation relative to bivalve ontogeny. *Journal of Shellfish Research*, 30(1), 187–196. <https://doi.org/10.2983/035.030.0126>
- Fisher, R. A., Call, G. C., & Grubbs, R. D. (2013). Age, growth, and reproductive biology of cownose rays in Chesapeake Bay. *Marine and Coastal Fisheries*, 5(1), 224–235. <https://doi.org/10.1080/19425120.2013.812587>
- Flach, E., & Tamaki, A. (2001). Competitive bioturbators on intertidal sand flats in the European Wadden Sea and Ariake Sound in Japan. In *Ecological Comparisons of Sedimentary Shores* (Vol. 151, pp. 149–171). <https://doi.org/10.1007/978-3-642-56557-18>
- Flowers, K. I., Heithaus, M. R., & Papastamatiou, Y. P. (2021). Buried in the sand: Uncovering the ecological roles and importance of rays. *Fish and Fisheries*, 22(1), 105–127. <https://doi.org/10.1111/faf.12508>
- Fokkema, W., Jeugd, H. P., Lameris, T. K., Dokter, A. M., Ebbinge, B. S., Roos, A. M., & Olf, H. (2020). Ontogenetic niche shifts as a driver of seasonal migration. *Oecologia*, 193(2), 285–297. <https://doi.org/10.1007/s00442-020-04682-0>
- Fox, H. E., Soltanoff, C. S., Mascia, M. B., Haisfield, K. M., Lombana, A. V., Pyke, C. R., & Wood, L. (2012). Explaining global patterns and trends in marine protected area (MPA) development. *Marine Policy*, 36(5), 1131–1138. <https://doi.org/10.1016/j.marpol.2012.02.007>
- Freitas, R. H. A., Aguiar, A. A., AKCHA, F., Lima, S. M. Q., & Valentin, J. L. (2019). Unravelling the foraging behavior of the southern stingray, *Hypanus americanus* (Myliobatiformes: Dasyatidae). In *In a southwestern Atlantic MPA. Neotrop ichthyol* 17. <http://www.scielo.br/j/ni/a/fZ57Gygh6b5nLVxrBsQy3b5/abstract/?lang=en>
- Friedman, K., Gabriel, S., Abe, O., Adnan Nuruddin, A., Ali, A., Bidin Raja Hassan, R., & Ye, Y. (2018). Examining the impact of CITES listing of sharks and rays in Southeast Asian fisheries. *Fish and Fisheries*, 19(4), 662–676. <https://doi.org/10.1111/faf.12281>
- Frisch, A. J., Ireland, M., Rizzari, J. R., Lönnstedt, O. M., Magnenat, K. A., Mirbach, C. E., & Hobbs, J. P. A. (2016). Reassessing the trophic role of reef sharks as apex predators on coral reefs. *Coral Reefs* (Online), 35(2), 459–472. <https://doi.org/10.1007/s00338-016-1415-2>

- Frisk, M. G., Miller, T. J., & Fogarty, M. J. (2001). Estimation and analysis of biological parameters in elasmobranch fishes: A comparative life history study. *Canadian Journal of Fisheries and Aquatic Sciences*, 58(5), 969–981. <https://doi.org/10.1139/f01-051>
- Froese, R. (2004). Keep it simple: Three indicators to deal with overfishing. *Fish and Fisheries*, 5(1), 86–91. <https://doi.org/10.1111/j.1467-2979.2004.00144.x>
- Frøsløv, T. G., Kjølner, R., Bruun, H. H., Ejrnæs, R., Brunbjerg, A. K., Pietroni, C., & Hansen, A. J. (2017). Algorithm for post-clustering curation of DNA amplicon data yields reliable biodiversity estimates. *Nature Communications*, 8(1). <https://doi.org/10.1038/s41467-017-01312-x>

G

- Gagern, A., & Bergh, J. (2013). A critical review of fishing agreements with tropical developing countries. *Marine Policy*, 38, 375–386. <https://doi.org/10.4324/9781351216227>
- Galbraith, H., Jones, R., Park, R., Clough, J., Herrod-Julius, S., Harrington, B., & Page, G. (2002). Global climate change and sea level rise: Potential losses of intertidal habitat for shorebirds. *Waterbirds*, 25(2), 173–183. [https://doi.org/10.1675/1524-4695\(2002\)025](https://doi.org/10.1675/1524-4695(2002)025)
- Gall, S. C., & Thompson, R. C. (2015). The impact of debris on marine life. *Marine Pollution Bulletin*, 92(1–2), 170–179. <https://doi.org/10.1016/j.marpolbul.2014.12.041>
- Gallagher, A. J., Amon, D. J., Bervoets, T., Shipley, O. N., Hammerschlag, N., & Sims, D. W. (2020). The Caribbean needs big marine protected areas. *Science (New York, N.Y.)*, 367(6479), 749. <https://doi.org/10.1126/science.abb0650>
- Garcia, S. M., Zerbi, A., Aliaume, C., Do Chi, T., & Lasserre, G. (2003). The ecosystem approach to fisheries. *FAO Fisheries Technical Paper*, 443.
- Garvey, J. E., & Whiles, M. (2016). *Trophic Ecology*. Trophic Ecology. CRC Press. <https://doi.org/10.1201/9781315367804>
- Gelsleichter, J., Musick, J. A., & Nichols, S. (1999). Food habits of the smooth dogfish, *Mustelus canis*, dusky shark, *Carcharhinus obscurus*, Atlantic sharpnose shark, *Rhizoprionodon terraenovae*, and the sand tiger, *Carcharias taurus*, from the northwest Atlantic Ocean. *Environmental Biology of Fishes*, 54, 205–217. <https://doi.org/10.1023/A>
- George, L. W., Martins, A. P. B., Heupel, M. R., & Simpfendorfer, C. A. (2019). Fine-scale movements of juvenile blacktip reef sharks *Carcharhinus melanopterus* in a shallow nearshore nursery. *Marine Ecology Progress Series*, 623, 85–97. <https://doi.org/10.3354/meps13010>
- Gervais, C. R., Nay, T. J., Renshaw, G., Johansen, J. L., Steffensen, J. F., & Rummer, J. L. (2018). Too hot to handle? Using movement to alleviate effects of elevated temperatures in a benthic elasmobranch. *Hemiscyllium Ocellatum*. *Marine Biology*, 165(11), 1–12. <https://doi.org/10.1007/s00227-018-3427-7>
- Gibson, R. N. (1986). Intertidal teleosts: Life in a fluctuating environment (pp. 388–408). *The Behaviour of Teleost Fishes*. <https://doi.org/10.1007/978-1-4684-8261-415>
- Gibson, R. N., & Yoshiyama, R. M. (1999). Intertidal fish communities. In M. H. Horn, K. L. M. Martin, & M. A. Chotkowski (Eds.), *Intertidal fishes: Life in two worlds* (pp. 264–297). Academic Press.
- Gilchrist, G., Mallory, M., & Merkel, F. (2005). Can local ecological knowledge contribute to wildlife management? Case studies of migratory birds. *Ecology and Society*, 10(1). <https://doi.org/10.5751/ES-01275-100120>
- Gill, D. A., Mascia, M. B., Ahmadi, G. N., Glew, L., Lester, S. E., Barnes, M., & Fox, H. E. (2017). Capacity shortfalls hinder the performance of marine protected areas globally. *Nature*, 543(7647), 665–669. <https://doi.org/10.1038/nature21708>

- Gils, J. A., Geest, M., Jansen, E. J., Govers, L. L., Fouw, J., & Piersma, T. (2012). Trophic cascade induced by molluscivore predator alters pore-water biogeochemistry via competitive release of prey. *Ecology*, 93(5), 1143–1152. <https://doi.org/10.1890/11-1282.1>
- Gils, J. A., Geest, M., Leyrer, J., Oudman, T., Lok, T., Onrust, J., & Piersma, T. (2013). Toxin constraint explains diet choice, survival and population dynamics in a molluscivore shorebird. *Proceedings of the Royal Society B: Biological Sciences*, 280(1763). <https://doi.org/10.1098/rspb.2013.0861>
- Gils, J. A., Lisovski, S., Lok, T., Meissner, W., Ozarowska, A., Fouw, J., & Klaassen, M. (2016). Climate change: Body shrinkage due to Arctic warming reduces red knot fitness in tropical wintering range. *Science (New York, N.Y.)*, 352(6287), 819–821. <https://doi.org/10.1126/science.aad6351>
- Gils, J. A., Piersma, T., Dekinga, A., & Dietz, M. W. (2003). Cost-benefit analysis of mollusc-eating in a shorebird: II. Optimizing gizzard size in the face of seasonal demands. *Journal of Experimental Biology*, 206(19), 3369–3380. <https://doi.org/10.1242/jeb.00546>
- Giorgini, M., Miguez, A., Esquius, K. S., Astarloa, C. D., Iribarne, O., Fanjul, E., & Escapa, M. (2019). Regenerative bioturbation by intertidal burrowing crabs modifies microphytobenthic composition and enhances primary production in SW Atlantic mudflats. *Marine Ecology Progress Series*, 632, 43–57. <https://doi.org/10.3354/meps13134>
- Glaspie, C. N., & Seitz, R. D. (2017). Role of habitat and predators in maintaining functional diversity of estuarine bivalves. *Marine Ecology Progress Series*, 570(April), 113–125. <https://doi.org/10.3354/meps12103>
- Glaus, K. B. J., Adrian-Kalchhauser, I., Piovano, S., Appleyard, S. A., Brunnschweiler, J. M., & Rico, C. (2019). Fishing for profit or food? Socio-economic drivers and fishers' attitudes towards sharks in Fiji. *Marine Policy*, 100(November 2018), 249–257. <https://doi.org/10.1016/j.marpol.2018.11.037>
- Golden, C. D., Allison, E. H., Dey, M. M., Halpern, B. S., J., M. D., Matthew, S., & Vaitla, B. (2016). Fall in fish catch threatens human health. *Nature*, 534(7607), 317–320.
- Goldenberg, S. B., & Shapiro, L. J. (1996). Physical mechanisms for the association of el niño and west african rainfall with atlantic major hurricane activity. *Journal of Climate*, 9, 1169–1187.
- Goodman, L. A. (1961). Snowball Sampling. *The Annals of Mathematical Statistics*, 32(1), 148–170. <https://doi.org/10.1214/aoms/1177705148>
- Goodwin, N. B., Dulvy, N. K., & Reynolds, J. D. (2002). Life-history correlates of the evolution of live bearing in fishes. *Philosophical Transactions of the Royal Society B: Biological Sciences*, 357(1419), 259–267. <https://doi.org/10.1098/rstb.2001.0958>
- Govers, L. L., Pieck, T., Bouma, T. J., Suykerbuyk, W., Smolders, A. J. P., & Katwijk, M. M. (2014). Seagrasses are negatively affected by organic matter loading and *Arenicola marina* activity in a laboratory experiment. *Oecologia*, 175(2), 677–685. <https://doi.org/10.1007/s00442-014-2916-8>
- Granadeiro, J. P., Belo, J., Henriques, M., Catalão, J., & Catry, T. (2021). Using sentinel-2 images to estimate topography, tidal-stage lags and exposure periods over large intertidal areas. *Remote Sensing*, 13, 1–17.
- Grant, J. (1983). The relative magnitude of biological and physical sediment reworking in an intertidal community. *Journal of Marine Research*, 41(4), 673–689. <https://doi.org/10.1357/002224083788520469>
- Gray, A. E., Mulligan, T. J., & Hannah, R. W. (1997). Food habits, occurrence, and population structure of the batray, *Myliobatis californica*, in Humboldt Bay, California. *Environmental Biology of Fishes*, 49(2), 227–238. <https://doi.org/10.1023/A:1007379606233>

- Grecian, W. J., Witt, M. J., Attrill, M. J., Bearhop, S., Becker, P. H., Egevang, C., & Votier, S. C. (2016). Seabird diversity hotspot linked to ocean productivity in the Canary Current Large Marine Ecosystem. *Biology Letters*, 12(8), 20160024. <https://doi.org/10.1098/rsbl.2016.0024>
- Griffen, B. D., & Byers, J. E. (2006). Intraguild predation reduces redundancy of predator species in multiple predator assemblage. *Journal of Animal Ecology*, 75(4), 959–966. <https://doi.org/10.1111/j.1365-2656.2006.01115.x>
- Grubbs, R. D., Carlson, J. K., Romine, J. G., Curtis, T. H., McElroy, W. D., McCandless, C. T., & Musick, J. A. (2016). Critical assessment and ramifications of a purported marine trophic cascade. *Scientific Reports*, 6(February), 20970. <https://doi.org/10.1038/srep20970>
- Guénette, S., Meissa, B., & Gascuel, D. (2014). Assessing the contribution of marine protected areas to the trophic functioning of ecosystems: A model for the Banc d'Arguin and the Mauritanian shelf. *PLoS ONE*, 9(4), 94742. <https://doi.org/10.1371/journal.pone.0094742>
- Guillemot, N., Rocklin, D., Chen, C., & Le, M. (2014). A framework for mapping small-scale coastal fisheries using fishers' knowledge. *ICES Journal of Marine Science*, 71, 1781–1792.

H

- Haeseker, S. L., & Cech, J. J. (1994). Food habits of the brown smoothhound shark (*Mustelus henlei*) from two sites in Tomales Bay. *California Fish and Game*, 79, 96–114.
- Halpern, B. S., Walbridge, S., Selkoe, K. A., Kappel, C. V., Micheli, F., D'Agrosa, C., & Watson, R. (2008). A global map of human impact on marine ecosystems. *Science (New York, N.Y.)*, 319(5865), 948–952. <https://doi.org/10.1126/science.1149345>
- Halpin, P. M. (2000). Habitat use by an intertidal salt-marsh fish: Trade-offs between predation and growth. *Marine Ecology Progress Series*, 198, 203–214. <https://doi.org/10.3354/meps198203>
- Hamady, L. L., Natanson, L. J., Skomal, G. B., & Thorrold, S. R. (2014). Vertebral bomb radiocarbon suggests extreme longevity in white sharks. *PLoS ONE*, 9(1), 84006. <https://doi.org/10.1371/journal.pone.0084006>
- Hammerschlag, N., Broderick, A. C., Coker, J. W., Coyne, M. S., Dodd, M., Frick, M. G., & Hawkes, L. A. (2015). Evaluating the landscape of fear between apex predatory sharks and mobile sea turtles across a large dynamic seascape. *Ecology*, 96(8), 2117–2126. <https://doi.org/10.1890/14-2113.1>
- Hammerschlag, N., Fallows, C., Meyer, M., Seakamela, S. M., Orndorff, S., Kirkman, S., & Creel, S. (2022). Loss of an apex predator in the wild induces physiological and behavioural changes in prey. *Biology Letters*, 18(1). <https://doi.org/10.1098/rsbl.2021.0476>
- Hammerschlag, N., Meyer, M., Seakamela, S. M., Kirkman, S., Fallows, C., & Creel, S. (2017). Physiological stress responses to natural variation in predation risk: Evidence from white sharks and seals. *Ecology*, 98(12), 3199–3210. <https://doi.org/10.1002/ecy.2049>
- Hammerschlag, N., Schmitz, O. J., Flecker, A. S., Lafferty, K. D., Sih, A., Atwood, T. B., & Cooke, S. J. (2019). Ecosystem function and services of aquatic predators in the anthropocene. *Trends in Ecology and Evolution*, 34(4), 369–383. <https://doi.org/10.1016/j.tree.2019.01.005>
- Hammerschlag, N., Williams, L., Fallows, M., & Fallows, C. (2019). Disappearance of white sharks leads to the novel emergence of an allopatric apex predator, the sevengill shark. *Scientific Reports*, 9(1), 6–11. <https://doi.org/10.1038/s41598-018-37576-6>
- Hanski, I. (1991). Specialist predators, Generalist predators and the Microtine Rodent cycle. *Journal of Applied Ecology*, 60(1), 353–367.

- Haque, A. B., Cavanagh, R. D., & Seddon, N. (2021). Evaluating artisanal fishing of globally threatened sharks and rays in the Bay of Bengal, Bangladesh. *PLoS ONE*, 16. <https://doi.org/10.1371/journal.pone.0256146>
- Haque, A. B., & Spaet, J. L. Y. (2021). Trade in threatened elasmobranchs in the Bay of Bengal, Bangladesh. *Fisheries Research*, 243(June), 106059. <https://doi.org/10.1016/j.fishres.2021.106059>
- Haque, A. B., Washim, M., D'Costa, N. G., Baroi, A. R., Hossain, N., Nanjiba, R., & Khan, N. A. (2021). Socio-ecological approach on the fishing and trade of rhino rays (Elasmobranchii: Rhinopristiformes) for their biological conservation in the Bay of Bengal, Bangladesh. *Ocean and Coastal Management*, 210(August 2020), 105690. <https://doi.org/10.1016/j.ocecoaman.2021.105690>
- Hardin, G. (1960). The competitive exclusion principle. *Science*, 131(3409), 1292–1297.
- Harms-Tuohy, C. A., Schizas, N. V., & Appeldoorn, R. S. (2016). Use of DNA metabarcoding for stomach content analysis in the invasive lionfish *Pterois volitans* in Puerto Rico. *Marine Ecology Progress Series*, 558, 181–191. <https://doi.org/10.3354/meps11738>.
- Hartig, F. (2023). DHARMA: Residual diagnostics for hierarchical (multi-level/mixed) regression models. <http://florianhartig.github.io/DHARMA/>
- Hasan, M. M., Nazru, K. M. S., Rashed Parvej, M., Alam Patwary, M. S., & Uddin, A. M. M. B. (2017). Shark and shark products trade channel and its conservation aspects in Bangladesh. *Journal of Fisheries & Livestock Production*, 05(01), 1–8. <https://doi.org/10.4172/2332-2608.1000221>
- Hauer, M. E., Fussell, E., Mueller, V., Burkett, M., Call, M., Abel, K., & Wrathall, D. (2020). Sea-level rise and human migration. *Nature Reviews Earth and Environment*, 1(1), 28–39. <https://doi.org/10.1038/s43017-019-0002-9>
- Hays, G. C., Ferreira, L. C., Sequeira, A. M. M., Meekan, M. G., Duarte, C. M., Bailey, H., & Thums, M. (2016). Key questions in marine megafauna movement ecology. *Trends in Ecology and Evolution*, 31(6), 463–475. <https://doi.org/10.1016/j.tree.2016.02.015>
- Hazen, E. L., Abrahms, B., Brodie, S., Carroll, G., Jacox, M. G., Savoca, M. S., & Bograd, S. J. (2019). Marine top predators as climate and ecosystem sentinels. *Frontiers in Ecology and the Environment*, 17(10), 565–574. <https://doi.org/10.1002/fee.2125>
- Hazin, F., Fischer, A., & Broadhurst, M. (2001). Aspects of reproductive biology of the scalloped hammerhead shark, *Sphyrna lewini*, off northeastern Brazil. *Environmental Biology of Fishes*, 61, 151–159.
- Heithaus, M. (2001). The biology of tiger sharks, *Galeocerdo cuvier*, in Shark Bay, Western Australia: Sex ratio, size distribution, diet, and seasonal changes in catch rates. *Environmental Biology of Fishes*, 2, 25–36.
- Heithaus, M. R., & Dill, L. M. (2002). Food availability and tiger shark predation risk influence. *Ecology*, 83(2), 1–12.
- Heithaus, M. R., Dunn, R. E., Farabaugh, N. F., Lester, E., Madin, E., Meekan, M., & Wirsing, A. J. (2022). Advances in our understanding of the ecological importance of sharks and their relatives. In J. C. Carrier, C. A. Simpfendorfer, M. R. Heithaus, & K. E. Yopak (Eds.), *Biology of sharks and their relatives* (3rd editio, pp. 486–522). CRC Press.
- Heithaus, M. R., Frid, A., Vaudo, J. J., Worm, B., & Wirsing, A. J. (2010). Unraveling the ecological importance of elasmobranchs. In J. C. Carrier, J. A. Musick, & M. R. Heithaus (Eds.), *Sharks and their relatives II: Biodiversity, adaptive physiology, and conservation* (pp. 1–748). <https://doi.org/10.1201/9781420080483>

- Heithaus, M. R., Frid, A., Wirsing, A. J., & Worm, B. (2008). Predicting ecological consequences of marine top predator declines. *Trends in Ecology and Evolution*, 23(4), 202–210. <https://doi.org/10.1016/j.tree.2008.01.003>
- Heithaus, M. R., Wirsing, A. J., Burkholder, D., Thomson, J., & Dill, L. M. (2009). Towards a predictive framework for predator risk effects: The interaction of landscape features and prey escape tactics. *Journal of Animal Ecology*, 78(3), 556–562. <https://doi.org/10.1111/j.1365-2656.2008.01512.x>
- Helfield, J. M., & Naiman, R. J. (2006). Keystone interactions: Salmon and bear in riparian forests of Alaska. *Ecosystems (New York, N.Y.)*, 9(2), 167–180. <https://doi.org/10.1007/s10021-004-0063-5>
- Hellin, J., & Meijer, M. (2006). Guidelines for Value Chain Analysis. FAO.
- Hemming, V., Burgman, M. A., Hanea, A. M., McBride, M. F., & Wintle, B. C. (2018). A practical guide to structured expert elicitation using the IDEA protocol. *Methods in Ecology and Evolution*, 9(1), 169–180. <https://doi.org/10.1111/2041-210X.12857>
- Henriques, M., Belo, S., JR, J. M., H. A., JA, P., T, D. T., & Roomen, M. (2022). The Bijagós Archipelago: A key area for waterbirds of the East Atlantic Flyway. In R. M. C. G. & C. O (Eds.), *East atlantic flyway assessment 2020: The status of coastal waterbird populations and their sites. Wadden sea flyway initiative* (pp. 81–93). Wetlands International, BirdLife International.
- Hensley, R. A., McCoid, M. J., & Luer, C. A. (1998). Litter Variation in the Atlantic Guitarfish (Rhinobatidae: *Rhinobatos lentiginosus*) with Comments on Distribution in the Gulf of Mexico. *The Southwestern Naturalist*, 43(4), 501–504.
- Hernández, C. E., Neill, P. E., Pulgar, J. M., Ojeda, F. P., & Bozinovic, F. (2002). Water temperature fluctuations and territoriality in the intertidal zone: Two possible explanations for the elevational distribution of body size in *Graus nigra*. *Journal of Fish Biology*, 61(2), 472–488. <https://doi.org/10.1006/jfbi.2002.2054>
- Heupel, M. R., Knip, D. M., Simpfendorfer, C. A., & Dulvy, N. K. (2014). Sizing up the ecological role of sharks as predators. *Marine Ecology Progress Series*, 495, 291–298. <https://doi.org/10.3354/meps10597>
- Heupel, M. R., Simpfendorfer, C. a, Espinoza, M., Smoothey, A. F., Tobin, A., & Peddemors, V. (2015). Conservation challenges of sharks with continental scale migrations. *Frontiers in Marine Science*, 2(February), 1–7. <https://doi.org/10.3389/fmars.2015.00012>
- Higuera-Rivas, J. E., Hoyos-Padilla, E. M., Elorriaga-Verplancken, F. R., Rosales-Nanduca, H., Rosenthal, R., & Urbán, R. J. (2023). Orcas (*orcinus orca*) use different strategies to prey on rays in the Gulf of California. *Aquatic Mammals*, 49(1), 7–18. <https://doi.org/10.1578/AM.49.1.2023.7>
- Hilborn, R., Amoroso, R. O., Anderson, C. M., Baum, J. K., Branch, T. A., Costello, C., & Ye, Y. (2020). Effective fisheries management instrumental in improving fish stock status. *Proceedings of the National Academy of Sciences of the United States of America*, 117(4), 2218–2224. <https://doi.org/10.1073/pnas.1909726116>
- Hill, N. K., Woodworth, B. K., Phinn, S. R., Murray, N. J., & Fuller, R. A. (2021). Global protected-area coverage and human pressure on tidal flats. *Conservation Biology*, 35(3), 933–943. <https://doi.org/10.1111/cobi.13638>
- Hind, E. J. (2015). A review of the past, the present, and the future of fishers' knowledge research: A challenge to established fisheries science. *ICES Journal of Marine Science*, 2(72), 341–358.
- Hines, A. H., Whitlatch, R. B., Thrush, S. F., Hewitt, J. E., Cummings, V. J., Dayton, P. K., & Legendre, P. (1997). Nonlinear foraging response of a large marine predator to benthic prey: Eagle ray pits and bivalves in a New Zealand sandflat. *Journal of Experimental Marine Biology and Ecology*, 216(1–2), 191–210. [https://doi.org/10.1016/S0022-0981\(97\)00096-8](https://doi.org/10.1016/S0022-0981(97)00096-8)

- Hockey, P. A. R., & Bosman, A. L. (1986). Man as an intertidal predator in transkei: Disturbance, community convergence and management of a natural food resource. *Oikos* (Copenhagen, Denmark), 46(1), 3. <https://doi.org/10.2307/3565373>
- Hoffmayer, E. R., & Parsons, G. R. (2003). Food habits of three shark species from the Mississippi Sound in the northern Gulf of Mexico. *Southeastern Naturalist*, 2(2), 271–280. [https://doi.org/10.1656/1528-7092\(2003\)002](https://doi.org/10.1656/1528-7092(2003)002)
- Hofstede, R. ter. (2001). Incidental catches of pelagic megafauna by the EU pelagic fleet in the Mauritanian Exclusive Economic Zone during the year. RIVO-Netherlands Institute for Fisheries Research.
- Hoggarth, D. D., Abeyasekera, S., Arthur, R. I., Beddington, J. R., Burn, R. W., Halls, A. S., & Welcomme, R. L. (2006). Stock assessment for fishery management, A frame guide to the stock assessment tools of the Fisheries Management Science Programme. FAO. Fisheries Technical Paper. <https://doi.org/10.1007/s13398-014-0173-7.2>
- Hollensead, L. D., Grubbs, R. D., Carlson, J. K., & Bethea, D. M. (2016). Analysis of fine-scale daily movement patterns of juvenile *Pristis pectinata* within a nursery habitat. *Aquatic Conservation: Marine and Freshwater Ecosystems*, 26(3), 492–505. <https://doi.org/10.1002/aqc.2556>
- Holt, R. D., & Huxel, G. R. (2007). Alternative prey and the dynamics of intraguild predation: Theoretical perspectives. *Ecology*, 88(11), 2706–2712. <https://doi.org/10.1890/06-1525.1>
- Honda, K., Nakamura, Y., Nakaoka, M., Uy, W. H., & Fortes, M. D. (2013). Habitat use by fishes in coral reefs, seagrass beds and mangrove habitats in the Philippines. *PLoS ONE*, 8(8), 1–10. <https://doi.org/10.1371/journal.pone.0065735>
- Honkoop, P. J. C., Berghuis, E. M., Holthuijsen, S. J., Lavaleye, M. S. S., & Piersma, T. (2008). Molluscan assemblages of seagrass-covered and bare intertidal flats on the Banc d'Arguin, Mauritania, in relation to characteristics of sediment and organic matter. *Journal of Sea Research*, 60(4), 255–263. <https://doi.org/10.1016/j.seares.2008.07.005>
- Hooker, S. K., Cañadas, A., Hyrenbach, K. D., Corrigan, C., Polovina, J. J., & Reeves, R. R. (2011). Making protected area networks effective for marine top predators. *Endangered Species Research*, 13(3), 203–218. <https://doi.org/10.3354/esr00322>
- Hopsch, S. B., Thorncroft, C. D., Hodges, K., & Aiyyer, A. (2007). West African storm tracks and their relationship to atlantic tropical cyclones. *Journal of Climate*, 20, 2468–2483.
- Hughes, T. P., Kerry, J. T., Baird, A. H., Connolly, S. R., Dietzel, A., Eakin, C. M., & Torda, G. (2018). Global warming transforms coral reef assemblages. *Nature*, 556(7702), 492–496.
- Humber, F., Andriamahaino, E. T., Beriziny, T., Botosoamananto, R., Godley, B. J., Gough, C., & Broderick, A. C. (2017). Assessing the small-scale shark fishery of Madagascar through community-based monitoring and knowledge. *Fisheries Research*, 186, 131–143. <https://doi.org/10.1016/j.fishres.2016.08.012>
- Hussey, N. E., Brush, J., McCarthy, I. D., & Fisk, A. T. (2010). $\delta^{15}\text{N}$ and $\delta^{13}\text{C}$ diet-tissue discrimination factors for large sharks under semi-controlled conditions. *Comparative Biochemistry and Physiology - A Molecular and Integrative Physiology*, 155(4), 445–453. <https://doi.org/10.1016/j.cbpa.2009.09.023>
- Hussey, N. E., MacNeil, M. A., Siple, M. C., Popp, B. N., Dudley, S. F. J., & Fisk, A. T. (2015). Expanded trophic complexity among large sharks. *Food Webs*, 4, 1–7. <https://doi.org/10.1016/j.fooweb.2015.04.002>
- Hylton, S., White, W. T., & Chin, A. (2017). The sharks and rays of the Solomon Islands: A synthesis of their biological diversity, values and conservation status. *Pacific Conservation Biology*, 23(4), 324–334. <https://doi.org/10.1071/PC17012>

Hyslop, E. J. (1980). Stomach contents analysis—A review of methods and their application. *Journal of Fish Biology*, 17(4), 411–429. <https://doi.org/10.1111/j.1095-8649.1980.tb02775.x>.

I

IBAP. (2012). Archipel Des Bijagós – Motom Moranghajogo. IBAP.

ICAR. (2019). ICAR-CMFRI hosts FAO global expert meeting on shark trade. ICAR. <https://www.icar.org.in/content/icar-cmfri-hosts-fao-global-expert-meeting-shark-trade>

Ip, Y. C. A., Chang, J. J. M., Lim, K. K. P., Jaafar, Z., Wainwright, B. J., & Huang, D. (2021). Seeing through sedimented waters: Environmental DNA reduces the phantom diversity of sharks and rays in turbid marine habitats. *BMC Ecology and Evolution*, 21(1), 1–14. <https://doi.org/10.1186/s12862-021-01895-6>

IPBES. (2019). Summary for policymakers of the global assessment report on biodiversity and ecosystem services. In *Intergovernmental science-policy platform on biodiversity and ecosystem services (Vol. 45)*. <https://zenodo.org/record/3553579#.YfmYTerMI2w>

IPCC. (2007). *Climate change 2007: Impacts, adaptation and vulnerability. Contribution of working group II to the fourth assessment report of the intergovernmental panel on climate change*. <https://doi.org/10.1016/B978-008044910-4.00250-9>

Ishii, K., & Fukui, M. (2001). Optimization of annealing temperature to reduce bias caused by a primer mismatch in multitemplate PCR. *Applied and Environmental Microbiology*, 67(8), 3753–3755. <https://doi.org/10.1128/AEM.67.8.3753-3755.2001>

Ismen, A. (2003). Age, growth, reproduction and food of common stingray (*Dasyatis pastinaca* L., 1758) in Iskenderun Bay, the eastern Mediterranean. *Fisheries Research*, 60(1), 169–176. [https://doi.org/10.1016/S0165-7836\(02\)00058-9](https://doi.org/10.1016/S0165-7836(02)00058-9).

Iwamatsu, S., Suzuki, A., & Sato, M. (2007). Nereidid polychaetes as the major diet of migratory shorebirds on the estuarine tidal flats at Fujimae-Higata in Japan. *Zoological Science*, 24(7), 676–685. <https://doi.org/10.2108/zsj.24.676>

J

Jabado, R. W., Al Ghais, S. M., Hamza, W., & Henderson, A. C. (2015). The shark fishery in the United Arab Emirates: An interview based approach to assess the status of sharks. *Aquatic Conservation: Marine and Freshwater Ecosystems*, 25(6), 800–816. <https://doi.org/10.1002/aqc.2477>

Jabado, R. W., Al Ghais, S. M., Hamza, W., Shivji, M. S., & Henderson, A. C. (2015). Shark diversity in the Arabian/Persian Gulf higher than previously thought: Insights based on species composition of shark landings in the United Arab Emirates. *Marine Biodiversity*, 45(4), 719–731. <https://doi.org/10.1007/s12526-014-0275-7>

Jabado R. W., Badji L., Chartrain E., Bruyne G., Derrick D., Dia M., & Williams A. B. (2021). *Fontitrygon margaritella*. In *The IUCN Red List of Threatened Species* (p. 161520). <https://doi.org/10.2305/IUCN.UK.2021-2.RLTS.T161520A124498844.en>

Jabado, R. W., Chartrain, E., Bruyne, G., Derrick, D., Diop, M., Doherty, P., Keith Diagne, L., Leurs, G. H. L., Metcalfe, K., Sayer, C., Seidu, I., Tamo, A., VanderWright, W. J., & Williams, A. B. (2021). *Fontitrygon ukpam*. *The IUCN Red List of Threatened Species*.

Jabado, R. W., Chartrain, E., Cliff, G., Silva, C., Bruyne, G., Derrick, D., & Winker, H. (2021). *Mustelus mustelus*, common smoothhound. *The IUCN Red List of Threatened Species*, 39358 124405881. <https://doi.org/10.1007/978-3-030-50032-0141>

- Jabado R.W, Pacoureau N., Diop M., Dia M., Ba A., Williams A.B., & Metcalfe K. (2021). Rhinobatos rhinobatos, Common Guitarfish. In The IUCN Red List of Threatened Species 2021: E.T63131A124461877. <https://www.iucnredlist.org/species/63131/124461877>
- Jackson, J. B. C., Kirby, M. X., Berger, W. H., Bjorndal, K. A., Botsford, L. W., Bourque, B. J., & Warner, R. R. (2001). Historical overfishing and the recent collapse of coastal ecosystems. *Science (New York, N.Y.)*, 293(5530), 629–637. <https://doi.org/10.1126/science.1059199>
- Jacobsen, I. P., & Bennett, M. B. (2013). A Comparative Analysis of Feeding and Trophic Level Ecology in Stingrays (Rajiformes; Myliobatoidei) and Electric Rays (Rajiformes: Torpedinoidei). *PLoS ONE*, 8(8), e71348. <https://doi.org/10.1371/journal.pone.0071348>
- Jager, Z. (1993). The distribution and abundance of young fish in the Banc d'Arguin, Mauritania. *Hydrobiologia*, 258, 185–196. <https://doi.org/10.1007/BF00006196>
- Jaiteh, V. F., Lindfield, S. J., Mangubhai, S., Warren, C., Fitzpatrick, B., & Loneragan, N. (2016). Higher abundance of marine predators and changes in fishers' behavior following spatial protection within the world's biggest shark fishery. *Frontiers in Marine Science*, 3(April), 1–15. <https://doi.org/10.3389/fmars.2016.00043>
- Jennings, S., Pinnegar, J. K., Polunin, N. V. C., & Boon, T. W. (2001). Weak cross-species relationships between body size and trophic level belie powerful size-based trophic structuring in fish communities. *The Journal of Animal Ecology*, 70, 934–944.
- Jeunen, G. J., Knapp, M., Spencer, H. G., Lamare, M. D., Taylor, H. R., Stat, M., & Gemmill, N. J. (2019). Environmental DNA (eDNA) metabarcoding reveals strong discrimination among diverse marine habitats connected by water movement. *Molecular Ecology Resources*, 19(2), 426–438. <https://doi.org/10.1111/1755-0998.12982>
- Jin, B., Fu, C., Zhong, J., Li, B., Chen, J., & Wu, J. (2007). Fish utilization of a salt marsh intertidal creek in the Yangtze River Estuary, China. *Estuarine, Coastal and Shelf Science*, 73(3–4), 844–852. <https://doi.org/10.1016/j.ecss.2007.03.025>
- Jing, K., Ma, Z., Li, B., Li, J., & Chen, J. (2007). Foraging strategies involved in habitat use of shorebirds at the intertidal area of Chongming Dongtan, China. *Ecological Research*, 22(4), 559–570. <https://doi.org/10.1007/s11284-006-0302-7>
- Jing, Z., Kai, J., Xiaojing, G., & Zhijun, M. (2007). Food supply in intertidal area for shorebirds during stopover at Chongming Dongtan.
- Jirik, K. E., & Lowe, C. G. (2012). An elasmobranch maternity ward: Female round stingrays *Urobatis halleri* use warm, restored estuarine habitat during gestation. *Journal of Fish Biology*, 80(5), 1227–1245. <https://doi.org/10.1111/j.1095-8649.2011.03208.x>
- Joung, S. J., Chen, C. T., Clark, E., Uchida, S., & Huang, W. Y. P. (1996). The whale shark, *Rhincodon typus*, is a livebearer: 300 embryos found in one “megamamma” supreme. *Environmental Biology of Fishes*, 46(3), 219–223. <https://doi.org/10.1007/BF00004997>
- Juan-Jordá, M. J., Murua, H., Arrizabalaga, H., Merino, G., Pacoureau, N., & Dulvy, N. (2022). Seventy years of tunas, billfishes. And Sharks as Sentinels of Global Ocean Health, 378(6620). <https://doi.org/10.1126/science.abj0211>

K

- Kaiser, M. J., Attrill, M. J., Jennings, S., Thomas, D. N., Barnes, D. K. A., Brierley, A. S., & Kaartokallio, H. (2020). *Marine ecology: Processes, systems, and impacts* (3rd ed.). Oxford University Press.
- Kajiura, S. M., Sebastian, A. P., & Tricas, T. C. (2000). Dermal Bite Wounds as Indicators of Reproductive Seasonality and Behaviour in the Atlantic Stingray, *Dasyatis sabina*. *Environmental Biology of Fishes*, 58(1), 23–31. <https://doi.org/10.1023/A:1007667108362>

- Kalejta, B. (1993). Intense predation cannot always be detected experimentally. *Netherlands Journal of Sea Research*, 31(4), 385–393.
- Kanno, S., Heupel, M., & Simpfendorfer, C. (2019). Stationary video monitoring reveals habitat use of stingrays in mangroves. *Marine Ecology Progress Series*, 621, 155–168. <https://doi.org/10.3354/meps12977>
- Karnad, D., Sutaria, D., & Jabado, R. W. (2020). Local drivers of declining shark fisheries in India. *Ambio*, 49(2), 616–627. <https://doi.org/10.1007/s13280-019-01203-z>
- Keith, D. A., Rodríguez, J. P., Brooks, T. M., Burgman, M. A., Barrow, E. G., Bland, L., & Spalding, M. D. (2015). The IUCN red list of ecosystems: Motivations, challenges, and applications. *Conservation Letters*, 8(3), 214–226. <https://doi.org/10.1111/conl.12167>
- Kelly, R. P., Gallego, R., & Jacobs-Palme, E. (2018). The effect of tides on nearshore environmental DNA. *PeerJ*, 2018(3). <https://doi.org/10.7717/peerj.4521>
- Kempster, R. M. (2013). Sexual dimorphism of the electrosensory system: A quantitative analysis of nerve axons in the dorsal anterior lateral line nerve of the blue-spotted fantail stingray (*Taeniura lymma*). *Brain, Behavior and Evolution*, 81(4), 226–235. <https://doi.org/10.1159/000351700>.
- Kernaléguen, L., Dorville, N., Ierodiaconou, D., Hoskins, A. J., Baylis, A. M. M., Hindell, M. A., & Arnould, J. P. Y. (2016). From video recordings to whisker stable isotopes: A critical evaluation of timescale in assessing individual foraging specialisation in Australian fur seals. *Oecologia*, 180(3), 657–670. <https://doi.org/10.1007/s00442-015-3407-2>
- Kessel, S. T., & Hussey, N. E. (2015). Tonic immobility as an anaesthetic for elasmobranchs during surgical implantation procedures. *Canadian Journal of Fisheries and Aquatic Sciences*, 72(March), 1287–1291. <https://doi.org/10.1139/cjfas-2015-0136>
- Kiljunen, M., Grey, J., Sinisalo, T., Harrod, C., Immonen, H., & Jones, R. I. (2006). A revised model for lipid-normalizing $\delta^{13}\text{C}$ values from aquatic organisms, with implications for isotope mixing models. *Journal of Applied Ecology*, 43(6), 1213–1222. <https://doi.org/10.1111/j.1365-2664.2006.01224.x>
- Kim, S. L., Casper, D. R., Galván-Magaña, F., Ochoa-Díaz, R., Hernández-Aguilar, S. B., & Koch, P. L. (2012). Carbon and nitrogen discrimination factors for elasmobranch soft tissues based on a long-term controlled feeding study. *Environmental Biology of Fishes*, 95(1), 37–52. <https://doi.org/10.1007/s10641-011-9919-7>
- Kindsvater, H. K., Mangel, M., Reynolds, J. D., & Dulvy, N. K. (2016). Ten principles from evolutionary ecology essential for effective marine conservation. *Ecology and Evolution*, 6(7), 2125–2138. <https://doi.org/10.1002/ece3.2012>
- Kinsey, D. W., & Kinsey, B. E. (1967). Diurnal changes in oxygen content of the water over the coral reef platform at heron island. *Australian Journal of Marine and Freshwater Research*, 18, 23–34.
- Kleiber, P., Clarke, S., Bigelow, K., Nakano, H., Mcallister, M., Takeuchi, Y., Leeney, R. H., & Poncet, P. (2009). North pacific blue shark stock assessment. U.S. Department of Commerce, NOAA Technical Memorandum NMFS-PIFSC-17, February), 9(3), 411–430. <https://doi.org/10.1002/aqc.2419>
- Klimley, A. P. (2013). *The biology of sharks and rays*. Univ. of Chicago Press.
- Klokov, K., Gerasimov, Y., & Syroechkovskiy, E. (2023). Assessment of hunting pressure on Arctic-nesting shorebirds: First results from the Northeast of Russia. *E3S Web of Conferences*, 378. <https://doi.org/10.1051/e3sconf/202337805003>

- Knip, D. M., Heupel, M. R., & Simpfendorfer, C. A. (2010). Sharks in nearshore environments: Models, importance, and consequences. *Marine Ecology Progress Series*, 402, 1–11. <https://doi.org/10.3354/meps08498>
- Knip, D. M., Heupel, M. R., Simpfendorfer, C. A., Tobin, A. J., & Moloney, J. (2011). Ontogenetic shifts in movement and habitat use of juvenile pigeye sharks *Carcharhinus amboinensis* in a tropical nearshore region. *Marine Ecology Progress Series*, 425, 233–246. <https://doi.org/10.3354/meps09006>
- Koch, E. W., Barbier, E. B., Silliman, B. R., Reed, D. J., Perillo, G. M. E., Hacker, S. D., & Wolanski, E. (2009). Non-linearity in ecosystem services: Temporal and spatial variability in coastal protection. *Frontiers in Ecology and the Environment*, 7(1), 29–37. <https://doi.org/10.1890/080126>
- Koch, R. (1999). The 80/20 principle: The secret of achieving more with less. Crown Currency. [https://doi.org/10.1016/s0024-6301\(97\)80978-8](https://doi.org/10.1016/s0024-6301(97)80978-8)
- Kohler, N. E., & Turner, P. A. (2001). Shark tagging: A review of conventional methods and studies. *Environmental Biology of Fishes*, 60(191–223), 191–223.
- Koppel J, Heide T, Altieri AH, Eriksson BK, Bouma TJ, Olff H, & Silliman BR. (2015). Long-distance interactions regulate the structure and resilience of coastal ecosystems. *Annual Review of Marine Science*, 7, 139–158.
- Krakstad, J., Olsen, M., & Wagúe, A. (2004). Survey of the pelagic fish resources off north west africa. FAO.
- Kristensen, E., & Kostka, J. E. (2013). Macrofaunal burrows and irrigation in marine sediment: Microbiological and biogeochemical interactions. In *Interactions between macro- and microorganisms in marine sediments* (pp. 125–157). <https://doi.org/10.1029/CE060p0125>
- Kristensen, E., Penha-Lopes, G., Delefosse, M., Valdemarsen, T., Quintana, C. O., & Banta, G. T. (2012). What is bioturbation? The need for a precise definition for fauna in aquatic sciences. *Marine Ecology Progress Series*, 446, 285–302. <https://doi.org/10.3354/meps09506>
- Kronfeld-Schor, N., & Dayan, T. (2003). Partitioning of time as an ecological resource. *Annual Review of Ecology, Evolution, and Systematics*, 34, 153–181. <https://doi.org/10.1146/annurev.ecolsys.34.011802.132435>
- Kroodsma, D. A., Mayorga, J., Hochberg, T., Miller, N. A., Boerder, K., Ferretti, F., & Worm, B. (2018). Tracking the global footprint of fisheries. *Science (New York, N.Y.)*, 359(6378), 904–908. <https://doi.org/10.1126/science.aao5646>
- Kruijssen, F., Tedesco, I., Ward, A., Pincus, L., Love, D., & Thorne-Lyman, A. L. (2020). Loss and waste in fish value chains: A review of the evidence from low and middle-income countries. *Global Food Security*, 26, 100434. <https://doi.org/10.1016/j.gfs.2020.100434>
- Kruskal, J., & Wish, M. (1978). *Multidimensional scaling*. SAGE Publications, Inc.
- Küpper, F. C., & Kamenos, N. A. (2018). The future of marine biodiversity and marine ecosystem functioning in UK coastal and territorial waters (including UK Overseas Territories). With an Emphasis on Marine Macrophyte Communities. *Botanica Marina*, 61(6), 521–535. <https://doi.org/10.1515/bot-2018-0076>
- Kuwae, T., Miyoshi, E., Hosokawa, S., Ichimi, K., Hosoya, J., Amano, T., & Elnor, R. W. (2012). Variable and complex food web structures revealed by exploring missing trophic links between birds and biofilm. *Ecology Letters*, 15(4), 347–356. <https://doi.org/10.1111/j.1461-0248.2012.01744.x>
- Kyalo, K. B., & Stephen, N. (2013). Shark bycatch—Small scale tuna fishery interactions along the Kenyan coast. Indian Ocean Tuna Commission. <https://aquadocs.org/bitstream/handle/1834/7747/ktf0455.pdf>

- Kyne, P. M., & Jabado, R. W. (2019). *Rhynchobatus luebberti*, African wedgefish. In The IUCN Red List of Threatened Species:2019, 8235.
- Kyne, P. M., Jabado, R. W., Rigby, C. L., Dharmadi, G., A., M., Pollock, C. M., & Dulvy, N. K. (2020). The thin edge of the wedge: Extremely high extinction risk in wedgefishes and giant guitarfishes. *Aquatic Conservation: Marine and Freshwater Ecosystems*, 30(7), 1337–1361. <https://doi.org/10.1002/aqc.3331>
- ## L
- Lafferty, K. D., Benesh, K. C., Mahon, A. R., Jerde, C. L., & Lowe, C. G. (2018). Detecting southern California's white sharks with environmental DNA. *Frontiers in Marine Science*, 5(OCT), 1–6. <https://doi.org/10.3389/fmars.2018.00355>
- Lafferty, K. D., Garcia-Vedrenne, A. E., McLaughlin, J. P., Childress, J. N., Morse, M. F., & Jerde, C. L. (2020). At Palmyra Atoll, the fish-community environmental DNA signal changes across habitats but not with tides. *Journal of Fish Biology*. <https://doi.org/10.1111/jfb.14403>
- Lambarri, C., Espinosa, H., Martínez, A., & Hernández, A. (2015). Cods for sale Do we know what we are buying? *DNA Barcodes*, 3(1). <https://doi.org/10.1515/dna-2015-0004>
- Lara-Mendoza, R. E., Márquez-Farías, J. F., & Román-Reyes, J. C. (2015). Feeding habits of the speckled guitarfish *Rhinobatos glaucostigma* (Elasmobranchii: Rhinobatidae). *Journal of Fish Biology*, 87(2), 311–322. <https://doi.org/10.1111/jfb.12720>
- Larson, W. A., Barry, P., Dokai, W., Maselko, J., Olson, J., & Baetscher, D. (2022). Leveraging eDNA metabarcoding to characterize nearshore fish communities in Southeast Alaska: Do habitat and tide matter? *Environmental DNA*, 1–13. <https://doi.org/10.1002/edn3.297>
- Last, P., White, W., Carvalho, M., Séret, B., Stehmann, M., & Naylor, G. (2016). *Rays of the World*. CSIRO Publishing.
- Lathuilière, C., Echevin, V., & Lévy, M. (2008). Seasonal and intraseasonal surface chlorophyll-a variability along the northwest African coast. *Journal of Geophysical Research: Oceans*, 113(5), 2000–2004. <https://doi.org/10.1029/2007JC004433>
- Laverock, B., Gilbert, J. A., Tait, K., Osborn, A. M., & Widdicombe, S. (2011). Bioturbation: Impact on the marine nitrogen cycle. *Biochemical Society Transactions*, 39(1), 315–320. <https://doi.org/10.1042/BST0390315>
- Layman, C. A., & Post, D. M. (2007). Can stable isotope ratios provide for community-wide measures of trophic structure? *Ecology*, 88(1), 42–48. <https://doi.org/10.1890/08-0167.1>
- Le Quesne, W. J. F., & Jennings, S. (2012). Predicting species vulnerability with minimal data to support rapid risk assessment of fishing impacts on biodiversity. *Journal of Applied Ecology*, 49(1), 20–28. <https://doi.org/10.1111/j.1365-2664.2011.02087.x>.
- Lear, K. O., Whitney, N. M., Morgan, D. L., Brewster, L. R., Whitty, J. M., Poulakis, G. R., & Gleiss, A. C. (2019). Thermal performance responses in free-ranging elasmobranchs depend on habitat use and body size. *Oecologia*, 191(4), 829–842. <https://doi.org/10.1007/s00442-019-04547-1>
- Lear, K. O., Whitney, N. M., Morris, J. J., & Gleiss, A. C. (2021). Temporal niche partitioning as a novel mechanism promoting co-existence of sympatric predators in marine systems. *Proceedings of the Royal Society B: Biological Sciences*, 288(1954). <https://doi.org/10.1098/rspb.2021.0816>
- Leeney, R. H., & Poncelet, P. (2015). Using fishers' ecological knowledge to assess the status and cultural importance of sawfish in Guinea-Bissau. *Aquatic Conservation: Marine and Freshwater Ecosystems*, 25(3), 411–430. <https://doi.org/10.1002/aqc.2419>

- Lehman, R. N., Poulakis, G. R., Scharer, R. M., Schweiss, K. E., Hendon, J. M., & Phillips, N. M. (2020). An environmental DNA tool for monitoring the status of the Critically Endangered Smalltooth Sawfish, *Pristis pectinata*, in the western Atlantic. *Conservation Genetics Resources*, 12(4), 621–629. <https://doi.org/10.1007/s12686-020-01149-5>
- Lemrabott, S. Y. (2023). Fish and fisheries dynamics at Banc d'Arguin, Mauritania [PhD. thesis, University of Groningen]. <https://doi.org/10.33612/diss.574910952>
- Lemrabott, S. Y. C., El-Hacen, E. M., Piersma, T., Sall, A. A., Sidina, E., Mahmoud, L. Y. A., & Leeuwen, A. (2023). Twenty years of monitoring reveal overfishing of bony fish stocks in the coastal national park Banc d'Arguin. In Mauritania. Aquatic conservation: Marine and freshwater ecosystems (pp. 833–844). <https://doi.org/10.1002/aqc.3948>
- Lemrabott, S. Y., Van Leeuwen, A., Piersma, T., Braham, C.-B., Ball, A. C., Araujo, A., Olf, H., & El-Hacen, E.-H. (2024). The chronology of overfishing in a remote West African coastal ecosystem. *Ecology and Society*, 29(1), art9. <https://doi.org/10.5751/ES-13902-290109>
- Lenth, R. (2019). emmeans: Estimated marginal means, aka least-squares means. <https://CRANR-proje>
- Lenton, T. M., Held, H., Kriegler, E., Hall, J. W., Lucht, W., Rahmstorf, S., & Schellnhuber, H. J. (2008). Tipping elements in the Earth's climate system. *Proceedings of the National Academy of Sciences of the United States of America*, 105(6), 1786–1793. <https://doi.org/10.1073/pnas.0705414105>
- Leurs, G., Nieuwenhuis, B. O., Zuidewind, T. J., Hijner, N., Olf, H., & Govers, L. L. (2023). Where land meets sea: Intertidal areas as key habitats for sharks and rays. *Fish and Fisheries*, 1–20. <https://doi.org/10.1111/faf.12735>
- Leurs, G., Reijden, K. J., Cheikhna Lemrabott, S. Y., Barry, I., Nonque, D. M., Olf, H., Ledo Pontes, S., Regalla, A., & Govers, L. L. (2021). Industrial fishing near west african marine protected areas and its potential effects on mobile marine predators. *Frontiers in Marine Science*, 8. <https://doi.org/10.3389/fmars.2021.602917>
- Leurs, G., Verkuil, Y. I., Hijner, N., Saalman, F., Dos Santos, L., Regalla, A., Ledo Pontes, S., Yang, L., Naylor, G. J. P., Olf, H., & Govers, L. L. (2023). Addressing data-deficiency of threatened sharks and rays in a highly dynamic coastal ecosystem using environmental DNA. *Ecological Indicators*, 154(110795).
- Lewis, T. L., Esler, D., & Boyd, W. S. (2007). Effects of predation by sea ducks on clam abundance in soft-bottom intertidal habitats. *Marine Ecology Progress Series*, 329, 131–144. <https://doi.org/10.3354/meps329131>
- Lewison, R. L., Crowder, L. B., Read, A. J., & Freeman, S. A. (2004). Understanding impacts of fisheries bycatch on marine megafauna. *Trends in Ecology and Evolution*, 19(11), 598–604. <https://doi.org/10.1016/j.tree.2004.09.004>
- Lewison, R. L., Crowder, L. B., Wallace, B. P., Moore, J. E., Cox, T., Zydelski, R., & Safina, C. (2014). Global patterns of marine mammal, seabird, and sea turtle bycatch reveal taxa-specific and cumulative megafauna hotspots. *Proceedings of the National Academy of Sciences*, 111(14), 5271–5276.
- Li, Y., Zhang, Y., Hussey, N. E., & Dai, X. (2016). Urea and lipid extraction treatment effects on $\delta^{15}\text{N}$ and $\delta^{13}\text{C}$ values in pelagic sharks. *Rapid Communications in Mass Spectrometry*, 30(1), 1–8. <https://doi.org/10.1002/rcm.7396>
- Lifjeld, J. T. (1984). Prey selection in relation to body size and bill length of five species of waders feeding in the same habitat. *Ornis Scandinavica*, 15(4), 217–226. <https://doi.org/10.2307/3675930>

- Lim, K. C., Chong, V. C., Lim, P.-E., Yurimoto, T., & Loh, K. H. (2019). Feeding ecology of three sympatric species of stingrays on a tropical mudflat. *Journal of the Marine Biological Association of the United Kingdom*, 99(4), 999–1007. <https://doi.org/10.1017/S0025315418000759>
- Lindegren, M., Holt, B. G., MacKenzie, B. R., & Rahbek, C. (2018). A global mismatch in the protection of multiple marine biodiversity components and ecosystem services. *Scientific Reports*, 8(1), 1–8. <https://doi.org/10.1038/s41598-018-22419-1>
- Logan, J. M., & Lutcavage, M. E. (2010). Stable isotope dynamics in elasmobranch fishes. *Hydrobiologia*, 644(1), 231–244. <https://doi.org/10.1007/s10750-010-0120-3>
- Lohrer, A. M., Thrush, S. F., & Gibbs, M. M. (2004). Bioturbators enhance ecosystem function through complex biogeochemical interactions. *Nature*, 431(7012), 1092–1095. <https://doi.org/10.1038/nature03042>
- Lonneville, B., Oset García, P., Schepers, L., Vanhoorne, B., Hernandez, F., & Mees, J. (2019). *MarineRegions.org*. <http://www.marinerregions.org>
- Lopes, P. F. M., Verba, J. T., Begossi, A., & Pennino, M. G. (2019). Predicting species distribution from fishers' local ecological knowledge: A new alternative for data-poor management. *Canadian Journal of Fisheries and Aquatic Sciences*, 76(8), 1423–1431. <https://doi.org/10.1139/cjfas-2018-0148>
- Loreau, M., Mouquet, N., & Holt, R. D. (2003). Meta-ecosystems: A theoretical framework for a spatial ecosystem ecology. *Ecology Letters*, 6(8), 673–679. <https://doi.org/10.1046/j.1461-0248.2003.00483.x>
- Lorenzo, M., Claudet, J., & Guidetti, P. (2016). Spillover from marine protected areas to adjacent fisheries has an ecological and a fishery component. *Journal for Nature Conservation*, 32, 62–66. <https://doi.org/10.1016/j.jnc.2016.04.004>
- Lotze, H. K. (2007). Rise and fall of fishing and marine resource use in the Wadden Sea, southern North Sea. *Fisheries Research*, 87(2–3), 208–218. <https://doi.org/10.1016/j.fishres.2006.12.009>
- Lotze, H. K., Lenihan, H. S., Bourque, B. J., Bradbury, R. H., Cooke, R. G., Kay, M. C., & Jackson, J. B. C. (2011). Depletion, degradation, and recovery potential of estuaries and coastal seas. *Journal of Physics A: Mathematical and Theoretical*, 44(8), 1689–1699. <https://doi.org/10.1088/1751-8113/44/8/085201>
- Lourenço, P. M., Catry, T., & Granadeiro, J. P. (2017). Diet and feeding ecology of the wintering shorebird assemblage in the Bijagós archipelago, Guinea-Bissau. *Journal of Sea Research*, 128(December 2016), 52–60. <https://doi.org/10.1016/j.seares.2017.08.004>
- Lourenço, P. M., Catry, T., Piersma, T., & Granadeiro, J. P. (2015). Comparative feeding ecology of shorebirds wintering at Banc d'Arguin, Mauritania. *Estuaries and Coasts*, 39(3), 855–865. <https://doi.org/10.1007/s12237-015-0029-1>
- Lourenço, P. M., Granadeiro, J. P., & Catry, T. (2018). Low macroinvertebrate biomass suggests limited food availability for shorebird communities in intertidal areas of the Bijagós archipelago (Guinea-Bissau). *Hydrobiologia*, 816, 197–212.
- Lu, Y., Yuan, J., Lu, X., Su, C., Zhang, Y., Wang, C., & Sweijd, N. (2018). Major threats of pollution and climate change to global coastal ecosystems and enhanced management for sustainability. *Environmental Pollution*, 239, 670–680. <https://doi.org/10.1016/j.envpol.2018.04.016>
- Lucifora, L. O., García, V. B., & Escalante, A. H. (2009). How can the feeding habits of the sand tiger shark influence the success of conservation programs? *Animal Conservation*, 12(4), 291–301. <https://doi.org/10.1111/j.1469-1795.2009.00247.x>

- Lucifora, L. O., García, V. B., Menni, R. C., & Escalante, A. H. (2006). Food habits, selectivity, and foraging modes of the school shark *Galeorhinus galeus*. *Marine Ecology Progress Series*, 315, 259–270. <https://doi.org/10.3354/meps315259>
- Lucifora, L. O., García, V. B., Menni, R. C., Escalante, A. H., & Hozbor, N. M. (2008). Effects of body size, age and maturity stage on diet in a large shark: Ecological and applied implications. *Ecological Research*, 24(1), 109–118. <https://doi.org/10.1007/s11284-008-0487-z>
- Lucifora, L. O., Menni, R. C., & Escalante, A. H. (2005). Reproduction, abundance and feeding habits of the broadnose sevengill shark *Notorynchus cepedianus* in north Patagonia, Argentina. *Marine Ecology Progress Series*, 289, 237–244.
- Lynn-Myrick, J., & Flessa, K. W. (1996). Bioturbation rates in Bahía la Choya, Sonora, Mexico. *Ciencias marinas*, 22, 23–46.
- Lyons, K. (2017). Who's My Daddy? Considerations for the influence of sexual selection on multiple paternity in elasmobranch mating systems. *Ecology and Evolution*, 7(15), 5603–5612. <https://doi.org/10.1002/ece3.3086>.

M

- Roomen M, Schekkerman H, Delany S, Winden E, Flink S, Langendoen T, & Nagy S. (2011). Overview of monitoring work on numbers, reproduction and survival of waterbird populations important in the Wadden Sea and the East Atlantic Flyway. SOVON. (2011).
- MacArthur, R. H., & Pianka, E. R. (1966). On optimal use of a patchy environment. *The American Naturalist*, 100(916), 603–609.
- MacKendrick, A. G., & Kesava, P. M. (1911). The rate of multiplication of micro-organisms: A mathematical study. *Proceedings of the Royal Society of Edinburgh*, 31, 649–653.
- MacKeracher, T., Diedrich, A., & Simpfendorfer, C. A. (2019). Sharks, rays and marine protected areas: A critical evaluation of current perspectives. *Fish and Fisheries*, 20(2), 255–267. <https://doi.org/10.1111/faf.12337>
- MacKinnon, J., Verkuil, Y. I., & Murray, N. (2012). IUCN situation analysis on East and Southeast Asian intertidal habitats, with particular reference to the Yellow Sea (including the Bohai Sea). *Occasional Paper of the IUCN Species Survival Commission*, 47.
- MacNeil, M. A., Drouillard, K. G., & Fisk, A. T. (2006). Variable uptake and elimination of stable nitrogen isotopes between tissues in fish. *Canadian Journal of Fisheries and Aquatic Sciences*, 63(2), 345–353. <https://doi.org/10.1139/f05-219>
- MacNeil, M. A., Skomal, G. B., & Fisk, A. T. (2005). Stable isotopes from multiple tissues reveal diet switching in sharks. *Marine Ecology Progress Series*, 302, 199–206. <https://doi.org/10.3354/meps302199>.
- Malpica-Cruz, L., Herzka, S. Z., Sosa-Nishizaki, O., Lazo, J. P., & Trudel, M. (2012). Tissue-specific isotope trophic discrimination factors and turnover rates in a marine elasmobranch: Empirical and modeling results. *Canadian Journal of Fisheries and Aquatic Sciences*, 69, 551–564. <https://doi.org/10.1139/f2011-172>
- Manjarrez Acosta, C., Juárez Rentería, F., Rodríguez Espinoza, J. P., Díaz Duran, R., Lizárraga Humaran, X., & Vega Cerecer, A. E. (1983). Estudio sobre algunos aspectos biológicos pesqueros del tiburón en la zona sur de Sinaloa. *Universidad Autónoma de Sinaloa*.
- Marshall L. J., & Cronin E. S. (2016). *Fontitrygon margaritella*. In *The IUCN Red List of Threatened Species 2016* (p. 161495 104172339, 8235).

- Marshall, L. J., White, W. T., & Potter, I. C. (2007). Reproductive biology and diet of the southern fiddler ray, *Trygonorrhina fasciata* (Batoidea: Rhinobatidae), an important trawl bycatch species. *Marine and Freshwater Research*, 58(1), 104–115. <https://doi.org/10.1071/MF05165>
- Martin, R. A. (2005). Conservation of freshwater and euryhaline elasmobranchs: A review. *Journal of the Marine Biological Association of the UK*, 85, 1049. <https://doi.org/10.1017/S0025315405012105>
- Martinell, J., Gibert, J. M., Domènech, R., Ekdale, A. A., & Steen, P. P. (2001). Cretaceous ray traces?: An alternative interpretation for the alleged dinosaur tracks of La Posa, Isona, NE Spain. *Palaios*, 16(4), 409–416.
- Martins, A. P. B., Heupel, M. R., Bierwagen, S. L., Chin, A., & Simpfendorfer, C. A. (2020). Tidal-diel patterns of movement, activity and habitat use by juvenile mangrove whiprays using towed-float GPS telemetry. *Marine and Freshwater Research*. <https://doi.org/10.1071/MF20078>
- Martins, A. P. B., Heupel, M. R., Chin, A., & Simpfendorfer, C. A. (2018). Batoid nurseries: Definition, use and importance. *Marine Ecology Progress Series*, 595, 253–267. <https://doi.org/10.3354/meps12545>
- Martins, A. P. B., Matley, J. K., Heupel, M. R., Fisk, A. T., Chin, A., & Simpfendorfer, C. A. (2022). Trophic ecology of sympatric juvenile stingrays within a nursery area. *Marine and Freshwater Research*, 1–11. <https://doi.org/10.1071/mf21292>
- Matern, S. A., Cech, J. J., & Hopkins, T. E. (2000). Diel movements of bat rays, *Myliobatis californica*, in Tomales Bay, California: Evidence for behavioral thermoregulation? *Environmental Biology of Fishes*, 58(2), 173–182. <https://doi.org/10.1023/A:1007625212099>
- Mathot, K., Piersma, T., & Elner, R. W. (2019). Shorebirds as Integrators and Indicators of Mudflat Ecology. In *Mudflat Ecology* (7th ed., pp. 309–338). Springer.
- Matich, P., Heithaus, M. R., & Layman, C. A. (2011). Contrasting patterns of individual specialization and trophic coupling in two marine apex predators. *Journal of Animal Ecology*, 80(1), 294–305. <https://doi.org/10.1111/j.1365-2656.2010.01753.x>
- Maunder, M. N., & Punt, A. E. (2004). Standardizing catch and effort data: A review of recent approaches. *Fish. Res.*, 70, 141–159. <https://doi.org/10.1016/j.fishres.2004.08.002>
- Maxwell, D., & Jennings, S. (2005). Power of monitoring programmes to detect decline and recovery of rare and vulnerable fish. *Journal of Applied Ecology*, 42(1), 25–37. <https://doi.org/10.1111/j.1365-2664.2005.01000.x>
- McCarter, J., & Gavin, M. C. (2014). Local perceptions of changes in traditional ecological knowledge: A case study from Malekula Island, Vanuatu. *Ambio*, 43(3), 288–296. <https://doi.org/10.1007/s13280-013-0431-5>
- McCauley, D. J., Young, H. S., Dunbar, R. B., Estes, J. A., Semmens, B. X., & Micheli, F. (2012). Assessing the effects of large mobile predators on ecosystem connectivity. *Ecological Applications*, 22(6), 1711–1717. <https://doi.org/10.1890/11-1653.1>
- McDavitt, M. T. (2005). The cultural significance of sharks and rays in Aboriginal societies across Australia's top end. *Seaweed*, 1–5.
- McDermott, G. R., Meng, K. C., McDonald, G. G., & Costello, C. J. (2018). The blue paradox: Preemptive overfishing in marine reserves. *Proceedings of the National Academy of Sciences*, 116(12), 5319–5325. <https://doi.org/10.1073/pnas.1802862115>
- Meijer, K. J., El-Hacen, E. H. M., Govers, L. L., Lavaleye, M., Piersma, T., & Olf, H. (2021). Mangrove-mudflat connectivity shapes benthic communities in a tropical intertidal system. *Ecological Indicators*, 130(December 2020), 108030. <https://doi.org/10.1016/j.ecolind.2021.108030>

- Mejia-Falla, P. A. (2014). Age and growth of the round stingray *Urotrygon rogersi*, a particularly fast-growing and short-lived elasmobranch. *PLoS ONE*, 9(4), 12–15. <https://doi.org/10.1371/journal.pone.0096077>.
- Mendonça, V. M., Raffaelli, D. G., & Boyle, P. R. (2007). Interactions between shorebirds and benthic invertebrates at Culbin Sands lagoon, NE Scotland: Effects of avian predation on their prey community density and structure. *Scientia Marina*, 71(3), 579–591. <https://doi.org/10.3989/scimar.2007.71n3579>
- Meredith, T. L., Kajiura, S. M., Newton, K. C., Tricas, T. C., & Bedore, C. N. (2022). Advances in the sensory biology of elasmobranchs. In J. C. Carrier, C. A. Simpfendorfer, M. R. Heithaus, & K. E. Yopak (Eds.), *Biology of sharks and their relatives* (third ed., pp. 143–168). CRC Press.
- Merem, E. C., Twumasi, Y., Wesley, J., Alsarari, M., Fageir, S., & Crisler, M. (2019). Analyzing the tragedy of illegal fishing on the west african coastal region. *Food Public Heal*, 9, 1–15. <https://doi.org/10.5923/j.food.20190901.01>.
- Merten, V., Christiansen, B., Javidpour, J., Piatkowski, U., Puebla, O., Gasca, R., & Hoving, H. J. T. (2017). Diet and stable isotope analyses reveal the feeding ecology of the orangeback squid *Sthenoteuthis pteropus* (Steenstrup 1855). In the Eastern Tropical Atlantic. *PLoS ONE*, 12(12). <https://doi.org/10.1371/journal.pone.0189691>
- Methot, R. D., & Wetzel, C. R. (2013). Stock synthesis: A biological and statistical framework for fish stock assessment and fishery management. *Fisheries Research*, 142, 86–99. <https://doi.org/10.1016/j.fishres.2012.10.012>
- Meyer, W., & Seegers, U. (2012). Basics of skin structure and function in elasmobranchs: A review. *Journal of Fish Biology*, 80(5), 1940–1967. <https://doi.org/10.1111/j.1095-8649.2011.03207.x>
- Meysman, F. J. R., Middelburg, J. J., & Heip, C. H. R. (2006). Bioturbation: A fresh look at Darwin's last idea. *Trends in Ecology and Evolution*, 21(12), 688–695. <https://doi.org/10.1016/j.tree.2006.08.002>
- Mills, L. S., & Doak, D. F. (1993). The keystone-species concept in ecology and conservation. *BioScience*, 43(4), 219–224. <https://doi.org/10.2307/1312122>
- Minamoto, T., Naka, T., Moji, K., & Maruyama, A. (2016). Techniques for the practical collection of environmental DNA: filter selection, preservation, and extraction. *Limnology*, 17(1), 23–32. <https://doi.org/10.1007/s10201-015-0457-4>
- Mitra, A., & Zaman, S. (2016). Threats to Marine and Estuarine Ecosystems. *Basics of Marine and Estuarine Ecology*. <https://doi.org/10.1007/978-81-322-2707-610>
- Miya, M. (2022). Environmental DNA metabarcoding: A novel method for biodiversity monitoring of marine fish communities. *Annual Review of Marine Science*, 14, 161–185. <https://doi.org/10.1146/annurev-marine-041421-082251>
- Mollet, H. F. (2000). Reproductive biology of the female shortfin mako, *Isurus oxyrinchus Rafinesque*, 1810, with comments on the embryonic development of lamnoids. *Fishery Bulletin*, 98(2), 299–318.
- Momigliano, P., & Harcourt, R. (2014). Sharks: Conservation, governance and management. In *Sharks: Conservation, governance and management* (pp. 89–106). <https://doi.org/10.4324/9780203750292>
- Moore, A. (2017). Guitarfishes: The next sawfishes? Extinction vulnerabilities and an urgent call for conservation action. *Endangered Species Research*, 34, 75–88. <https://doi.org/10.3354/esr00830>
- Moore, A. B. M., Séret, B., & Armstrong, R. (2019). Risks to biodiversity and coastal livelihoods from artisanal elasmobranch fisheries in a least developed country: The Gambia (West Africa) biodiversity and conservation. Springer Netherlands. <https://doi.org/10.1007/s10531-019-01732-9>.

- Moradi, S., Sheykhi Ilanloo, S., Kafash, A., & Yousefi, M. (2019). Identifying high-priority conservation areas for avian biodiversity using species distribution modeling. *Ecological Indicators*, 97(October 2018), 159–164. <https://doi.org/10.1016/j.ecolind.2018.10.003>
- Morrissey, J. F., & Gruber, S. H. (1993). Habitat selection by juvenile lemon sharks, *Negaprion brevirostris*. *Environmental Biology of Fishes*, 38(4), 311–319. <https://doi.org/10.1007/BF00007524>
- Moss, B. (2017). Marine reptiles, birds and mammals and nutrient transfers among the seas and the land: An appraisal of current knowledge. *Journal of Experimental Marine Biology and Ecology*, 492, 63–80. <https://doi.org/10.1016/j.jembe.2017.01.018>
- Moura, T., Figueiredo, I., Farias, I., Serra-Pereira, B., Neves, A., Borges, M. D. F., & Serrano Gordo, L. (2008). Ontogenetic dietary shift and feeding strategy of *Raja undulata* Lacepède, 1802 (Chondrichthyes: Rajidae) on the Portuguese continental shelf. *Scientia Marina*, 72(2), 311–318. <https://doi.org/10.3989/scimar.2008.72n2311>
- Mourier, J., & Planes, S. (2013). Direct genetic evidence for reproductive philopatry and associated fine-scale migrations in female blacktip reef sharks (*Carcharhinus melanopterus*) in French Polynesia. *Molecular Ecology*, 22(1), 201–214. <https://doi.org/10.1111/mec.12103>
- Mull, C. G., Andrzejczek, S., Udyawer, V., & Dwyer, R. G. (2022). Advances in methods, understanding, and applications of elasmobranch movement ecology. In *Biology of sharks and their relatives* (pp. 357–399). CRC Press.
- Munroe, S. E. M., Simpfendorfer, C. A., & Heupel, M. R. (2014). Habitat and space use of an abundant nearshore shark, *Rhizoprionodon taylori*. *Marine and Freshwater Research*, 65(11), 959–968. <https://doi.org/10.1071/MF13272>
- Murray, N. J., Phinn, S. R., DeWitt, M., Ferrari, R., Johnston, R., Lyons, M. B., & Fuller, R. A. (2019). The global distribution and trajectory of tidal flats. *Nature*, 565(7738), 222–225. <https://doi.org/10.1038/s41586-018-0805-8>
- Murray, N. J., Worthington, T. A., Bunting, P., Duce, S., Hagger, V., Lovelock, C. E., & Lyons, M. B. (2022). High-resolution mapping of losses and gains of Earth's tidal wetlands. *Science*, 376(6594), 744–749. <https://doi.org/10.1126/science.abm9583>
- Musa, S. M., Ripley, D. M., Moritz, T., & Shiels, H. A. (2020). Ocean warming and hypoxia affect embryonic growth, fitness and survival of small-spotted catsharks, *Scyliorhinus canicula*. *Journal of Fish Biology*, 97(1), 257–264. <https://doi.org/10.1111/jfb.14370>
- Muttaqin, E., Abdullah, A., Nurilmala, M., Ichsan, M., Simeone, B. M., Yulianto, I., & Booth, H. (2019). DNA-barcoding as molecular marker for seafood forensics: Species identification of locally consumed shark fish products in the world's largest shark fishery. *IOP Conference Series: Earth and Environmental Science*, 278(1). <https://doi.org/10.1088/1755-1315/278/1/012049>
- Myers, R. A., Baum, J. K., Shepherd, T. D., Powers, S. P., & Peterson, C. H. (2007). Cascading effects of the loss of apex predatory sharks from a coastal ocean. *Science (New York, N.Y.)*, 315(5820), 1846–1850. <https://doi.org/10.1126/science.1138657>

N

- Nagelkerken, I., Blaber, S. J. M., Bouillon, S., Green, P., Haywood, M., Kirton, L. G., & Somerfield, P. J. (2008). The habitat function of mangroves for terrestrial and marine fauna: A review. *Aquatic Botany*, 89(2), 155–185. <https://doi.org/10.1016/j.aquabot.2007.12.007>

- Nagelkerken, I., Velde, G., Gorissen, M. W., Meijer, G. J., Van't Hof, T., & Hartog, C. (2000). Importance of mangroves, seagrass beds and the shallow coral reef as a nursery for important coral reef fishes, using a visual census technique. *Estuarine, Coastal and Shelf Science*, 51(1), 31–44. <https://doi.org/10.1006/ecss.2000.0617>
- Navia, A. F., Cortés, E., & Mejía-Falla, P. A. (2010). Topological analysis of the ecological importance of elasmobranch fishes: A food web study on the Gulf of Tortugas, Colombia. *Ecological Modelling*, 221(24), 2918–2926. <https://doi.org/10.1016/j.ecolmodel.2010.09.006>
- Navia, A. F., Mejía-Falla, P. A., & Giraldo, A. (2007). Feeding ecology of elasmobranch fishes in coastal waters of the Colombian Eastern Tropical Pacific. *BMC Ecology*, 7, 2–11. <https://doi.org/10.1186/1472-6785-7-8>.
- Navia, A. F., Mejía-Falla, P. A., López-García, J., Giraldo, A., & Cruz-Escalona, V. H. (2016). How many trophic roles can elasmobranchs play in a marine tropical network? *Marine and Freshwater Research*, 68(7), 1342–1353. <https://doi.org/10.1071/MF16161>
- Naylor, G. J. P. (2012). A DNA sequence-based approach to the identification of shark and ray species and its implications for global elasmobranch diversity and parasitology. *Bulletin of the American Museum of Natural History*, 367. <https://doi.org/10.1206/754.1>.
- Naylor, G. J. P., Caira, J. N., Jensen, K., Rosana, K. A. M., Straube, N., & Lakner, C. (2012). Elasmobranch phylogeny: A mitochondrial estimate based on 595 species. *Biology of Sharks and Their Relatives*, 47–72. <https://doi.org/10.1201/b11867-9>
- Naylor, G. J. P., Ryburn, J. A., Fedrigo, O., & Lopez, A. (2005). Phylogenetic relationships among the major lineages of modern elasmobranchs. In W. C. Hamlett (Ed.), *Reproductive biology and phylogeny of chondrichthyes* (pp. 1–26). CRC Press.
- Needham, H. R., Pilditch, C. A., Lohrer, A. M., & Thrush, S. F. (2011). Context-specific bioturbation mediates changes to ecosystem functioning. *Ecosystems (New York, N.Y.)*, 14, 1096–1109.
- Neis, B., Schneider, D. C., Felt, L., Haedrich, R. L., Fischer, J., & Hutchings, J. A. (1999). Fisheries assessment: What can be learned from interviewing resource users? *Canadian Journal of Fisheries and Aquatic Sciences*, 56(10), 1949–1963. <https://doi.org/10.1139/f99-115>
- Neuert, C., Meitinger, K., Behr, D., & Schonlau, M. (2021). The use of open-ended questions in surveys. *A Journal for Quantitative Methods and Survey Methodology*, 15(1), 3–6.
- Newman, S. P., Handy, R. D., & Gruber, S. H. (2010). Diet and prey preference of juvenile lemon sharks *Negaprion brevirostris*. *Marine Ecology Progress Series*, 398, 221–234. <https://doi.org/10.3354/meps08334>
- Newsome, S. D., Rio, C., Bearhop, S., & Philips, D. L. (2007). A niche for isotopic ecology. *Frontiers in Ecology and the Environment*, 5(8), 429–436. <https://doi.org/10.1890/060150.01>
- Ngwenya, B. N., Mosepele, K. K., & Magole, L. (2012). A case for gender equity in governance of the Okavango Delta fisheries in Botswana. *Natural Resources Forum*, 36(2), 109–122. <https://doi.org/10.1111/j.1477-8947.2012.001450.x>
- Niedemüller, S., Ainsworth, G., Juan, S., Garcia, R., Ospina-Alvarez, A., Pita, P., & Villasante, S. (2021). The shark and ray meat network. WWF. <https://sharks.panda.org/images/downloads/392/WWFMMIGlobalsharkraymeattradereport2021lowres.pdf>
- Niella, Y., Raoult, V., Gaston, T., Goodman, K., Harcourt, R., Peddemors, V., & Smoothey, A. F. (2022). Reliance of young sharks on threatened estuarine habitats for nutrition implies susceptibility to climate change. *Estuarine, Coastal and Shelf Science*, 268(February), 107790. <https://doi.org/10.1016/j.ecss.2022.107790>

- Nielsen, J., Hedeholm, R. B., Heinemeier, J., Bushnell, P. G., Christiansen, J. S., Olsen, J., & Stuiver, M. (2016). Eye lens radiocarbon reveals centuries of longevity in the Greenland shark (*Somniosus microcephalus*). *Science*, 353(6300), 702–704. <https://doi.org/10.1126/science.aaf1703>
- Nifong, J. C., & Lowers, R. H. (2017). Reciprocal intraguild predation between Alligator mississippiensis (american alligator) and elasmobranchii in the southeastern United States. *Southeastern Naturalist*, 16(3), 383–397. <https://doi.org/10.1656/058.016.0306>
- Nilsson, G. E., & Östlund-Nilsson, S. (2006). Hypoxia tolerance in coral reef fishes. In A. L. Val, V. M. F. Almeida-Val, & D. J. Randall (Eds.), *The physiology of tropical fishes* (pp. 583–596). Academic Press.
- Nordell, S. E. (1994). Observations of the mating behavior and dentition of the round stingray, *Urolophus halleri*. *Environmental Biology of Fishes*, 39(3), 219–229. <https://doi.org/10.1007/BF00005124>.
- Nosal, A. P., Cartamil, D. P., Ammann, A. J., Bellquist, L. F., Ben-Aderet, N. J., Blincow, K. M., & Hastings, P. A. (2021). Triennial migration and philopatry in the critically endangered soupfin shark *Galeorhinus galeus*. *Journal of Applied Ecology*, 58(8), 1570–1582. <https://doi.org/10.1111/1365-2664.13848>
- Nowicki, R., Heithaus, M., Thomson, J., Burkholder, D., Gastrich, K., & Wirsing, A. (2019). Indirect legacy effects of an extreme climatic event on a marine megafaunal community. *Ecological Monographs*, 89(3), 1–20. <https://doi.org/10.1002/ecm.1365>

O

- OBIS. (2023). Ocean Biodiversity Information System. Intergovernmental Oceanographic Commission of UNESCO. OBIS. www.obis.org.
- Ogburn, M. B., Bangley, C. W., Aguilar, R., Fisher, R. A., Curran, M. C., Webb, S. F., & Hines, A. H. (2018). Migratory connectivity and philopatry of cownose rays *Rhinoptera bonasus* along the Atlantic coast, USA. *Marine Ecology Progress Series*, 602(Mayr 1963), 197–211. <https://doi.org/10.3354/meps12686>
- Oksanen, J. (2020). VEGAN: Community ecology package. <https://cran.r-project.org/package=vegan>.
- Oliver, S., Braccini, M., Newman, S. J., & Harvey, E. S. (2015). Global patterns in the bycatch of sharks and rays. *Marine Policy*, 54, 86–97. <https://doi.org/10.1016/j.marpol.2014.12.017>
- Ortega-ortiz, C. D., Lazcano-pacheco, C., Llamas-, M., Meza-yáñez, R., Ruano-cobian, S., & López-luna, D. G. (2023). Expanding information on the prey items and hunting tactics of the eastern tropical Pacific killer whale (*Orcinus orca*) ecotype. *Examines Mar Biol Oceanogr.*, 6(2), 1–8. <https://doi.org/10.31031/EIMBO.2023.06.000635>
- Osaer, F. (2015). Sexual development and maturity scale for the angel shark *Squatina squatina* (Elasmobranchii: Squatinidae), with comments on the adequacy of general maturity scales. *Sexuality and Early Development in Aquatic Organisms*, 1(2), 117–132. <https://doi.org/10.3354/sedao00012>.
- O'Shea, O. R. (2012). *The Ecology and Biology of Stingrays (Dasyatidae) at Ningaloo Reef* [Murdoch University]. <https://researchportal.murdoch.edu.au/esploro/outputs/doctoral/The-ecology-and-biology-of-stingrays/991005543838807891#file-1>
- O'Shea, O. R. (2013). Growth of tropical dasyatid rays estimated using a multi-analytical approach. *PLoS ONE*, 8(10), 1–8. <https://doi.org/10.1371/journal.pone.0077194>.

- O'Shea, O. R., Thums, M., Keulen, M., Kempster, R. M., & Meekan, M. G. (2013). Dietary partitioning by five sympatric species of stingray (Dasyatidae) on coral reefs. *Journal of Fish Biology*, 82, 1805–1820.
- O'Shea, O. R., Thums, M., Keulen, M., & Meekan, M. (2012). Bioturbation by stingrays at Ningaloo Reef, Western Australia. *Marine and Freshwater Research*, 63(3), 189–197. <https://doi.org/10.1071/MF11180>
- Oudman, T., Onrust, J., Fouw, J., Spaans, B., Piersma, T., & Gils, J. A. (2014). Digestive capacity and toxicity cause mixed diets in red knots that maximize energy intake rate. *American Naturalist*, 183(5), 650–659. <https://doi.org/10.1086/675759>
- Oudman, T., Schekkerman, H., Kidee, A., Roomen, M., Camara, M., Smit, C., & El-Hacen, E. H. M. (2020). Changes in the waterbird community of the Parc National du Banc d'Arguin. *Bird Conservation International*, 30(4), 618–633. <https://doi.org/10.1017/S0959270919000431>
- P**
- Pacoureau, N., Rigby, C. L., Kyne, P. M., Sherley, R. B., Winker, H., Carlson, J. K., T., S., H., K., A., V. R., M., C., M., S., C., E.-H., & M, E. H. (2021). Half a century of global decline in oceanic sharks and rays. *Nature*, 589(4), 567–571. <https://doi.org/10.1038/s41586-020-03173-9>. Oudman
- Page, G., & Whitacre, D. F. (1975). Raptor predation on wintering shorebirds. *The Condor*, 77, 73–83.
- Paine, R. T. (1974). Intertidal community structure—Experimental studies on the relationship between a dominant competitor and its principal predator. *Oecologia*, 15(2), 93–120. <https://doi.org/10.1007/BF00345739>
- Palomares, M. L. D., & Pauly, D. (2019). Coastal fisheries: The past, present, and possible futures. *Coasts and estuaries: The future*. Elsevier Inc. <https://doi.org/10.1016/B978-0-12-814003-1.00032-0>
- Palumbi, S. R., Sandifer, P. A., Allan, J. D., Beck, M. W., Fautin, D. G., Fogarty, M. J., & Wall, D. H. (2009). Managing for ocean biodiversity to sustain marine ecosystem services. *Frontiers in Ecology and the Environment*, 7(4), 204–211. <https://doi.org/10.1890/070135>
- Papastamatiou, Y. P., Watanabe, Y. Y., Bradley, D., Dee, L. E., Weng, K., Lowe, C. G., & Caselle, J. E. (2015). Drivers of daily routines in an ectothermic marine predator: Hunt warm, rest warmer? *PLoS ONE*, 10(6), 1–16. <https://doi.org/10.1371/journal.pone.0127807>
- Parker, C., Cranford, M., Oakes, N., & Leggett, M. (2012). The little biodiversity finance book: A guide to proactive investment in natural capital. CBD. https://www.cbd.int/financial/hlp/doc/literature/LittleBiodiversityFinanceBook_3rd%20edition.pdf
- Parton, K. J., Galloway, T. S., & Godley, B. J. (2019). Global review of shark and ray entanglement in anthropogenic marine debris. *Endanger. Species Res*, 39, 173–190. <https://doi.org/10.3354/esr00964>.
- Pauly, D. (1995). Anecdotes and the shifting baseline syndrome of fisheries. *Trends in Ecology & Evolution*, 10(10), 430. [https://doi.org/10.1016/S0169-5347\(00\)89171-5](https://doi.org/10.1016/S0169-5347(00)89171-5)
- Pauly, D., & Christensen, V. (1995). Primary production required to sustain global fisheries. *Nature*, 374.
- Pauly, D., Christensen, V., Dalsgaard, J., Froese, R., Pauly, D., Christensen, V., Jr, & T, F. (1998). Fishing Down Marine Food Webs. *Science (New York, N.Y.)*, 279(5352), 860–863.
- Peacor, S. D., Barton, B. T., Kimbro, D. L., Sih, A., & Sheriff, M. J. (2020). A framework and standardized terminology to facilitate the study of predation-risk effects. *Ecology*, 101(12), 1–10. <https://doi.org/10.1002/ecy.3152>

- Pendleton, L., Donato, D. C., Murray, B. C., Crooks, S., Jenkins, W. A., Sifleet, S., Craft, C., Fourqurean, J. W., Kauffman, J. B., Marbà, N., Megonigal, P., Pidgeon, E., Herr, D., Gordon, D., & Baldera, A. (2012). Estimating Global “Blue Carbon” Emissions from Conversion and Degradation of Vegetated Coastal Ecosystems. *PLoS ONE*, 7(9), e43542. <https://doi.org/10.1371/journal.pone.0043542>
- Penning, E., Verkuil, Y. I., Klunder, L., & Reneerkens, J. (2022). Sanderlings feed on a diverse spectrum of prey worldwide but primarily rely on brown shrimp in the Wadden Sea. *Ardea*, 110(2), 187–199. <https://doi.org/10.5253/arde.2022.a11>
- Peterson, C. H. (1991). Intertidal zonation of marine invertebrates in sand and mud. *American Scientist*, 79(3), 236–249.
- Peterson, C. H., Fodrie, J. F., Summerson, H. C., & Powers, S. P. (2001). Site-specific and density-dependent extinction of prey by schooling rays: Generation of a population sink in top-quality habitat for bay scallops. *Oecologia*, 129(3), 349–356. <https://doi.org/10.1007/s004420100742>
- Philippe, A. S., Pinaud, D., Cayatte, M. L., Goulevant, C., Lachaussée, N., Pineau, P., & Bocher, P. (2016). Influence of environmental gradients on the distribution of benthic resources available for shorebirds on intertidal mudflats of Yves Bay, France. *Estuarine, Coastal and Shelf Science*, 174, 71–81. <https://doi.org/10.1016/j.ecss.2016.03.013>
- Phillips, D. L., Inger, R., Bearhop, S., Jackson, A. L., Moore, J. W., Parnell, A. C., & Ward, E. J. (2014). Best practices for use of stable isotope mixing models in food-web studies. *Canadian Journal of Zoology*, 92(10), 823–835. <https://doi.org/10.1139/cjz-2014-0127>
- Pierce, S. J., & Bennett, M. B. (2010). Destined to decline? Intrinsic susceptibility of the threatened estuary stingray to anthropogenic impacts. *Marine and Freshwater Research*, 61(12), 1468–1481. <https://doi.org/10.1071/MF10073>
- Pierce, S. J., Scott-Holland, T. B., & Bennett, M. B. (2011). Community composition of elasmobranch fishes utilizing intertidal sand flats in Moreton Bay, Queensland, Australia. *Pacific Science*, 65(2), 235–247. <https://doi.org/10.2984/65.2.235>
- Piersma, T. (1987). Production by intertidal benthic animals and limits to their predation by shorebirds: A heuristic model. *Marine Ecology Progress Series*, 38(Smit 1981), 187–196. <https://doi.org/10.3354/meps038187>
- Piersma, T. (2012). What is habitat quality? Dissecting a research portfolio on shorebirds. In *Birds and habitat* (pp. 383–407). <https://doi.org/10.1017/cbo9781139021654.019>
- Piersma, T., Goeij, P., & Tulp, I. (1993). An evaluation of intertidal feeding habitats from a shorebird perspective: Towards relevant comparisons between temperate and tropical mudflats. *Netherlands Journal of Sea Research*, 31(4), 503–512. [https://doi.org/10.1016/0077-7579\(93\)90062-W](https://doi.org/10.1016/0077-7579(93)90062-W)
- Pimiento, C., Leprieur, F., Silvestro, D., Lefcheck, J. S., Albouy, C., Rasher, D. B., & Griffin, J. N. (2020). Functional diversity of marine megafauna in the Anthropocene. *Science Advances*, 6(16). <https://doi.org/10.1126/sciadv.aay7650>
- Pinnegar, J. K., Polunin, N. V. C., Francour, P., Badalamenti, F., Chemello, R., Harmelin-Vivien, M.-L., Hereu, B., Milazzo, M., Zabala, M., D’anna, G., & Pipitone, C. (2000). Trophic cascades in benthic marine ecosystems: Lessons for fisheries and protected-area management. *Environmental Conservation*, 27, 179–200.
- Platell, M. E., Potter, I. C., & Clarke, K. R. (1998). Resource partitioning by four species of elasmobranchs (Batoidea: Urolophidae) in coastal waters of temperate Australia. *Marine Biology*, 131(4), 719–734. <https://doi.org/10.1007/s002270050363>

- Polanco Fernández, A., Marques, V., Fopp, F., Juhel, J. B., Borrero-Pérez, G. H., Cheutin, M. C., & Pellissier, L. (2021). Comparing environmental DNA metabarcoding and underwater visual census to monitor tropical reef fishes. *Environmental DNA*, 3(1), 142–156. <https://doi.org/10.1002/edn3.140>
- Polunin, N. V. C., & Roberts, C. M. (1993). Greater biomass and value of target coral-reef fishes in two small Caribbean marine reserves. *Marine Ecology Progress Series*, 100(1–2), 167–176. <https://doi.org/10.3354/meps100167>
- Pomeroy, R. S. (2012). Managing overcapacity in small-scale fisheries in Southeast Asia. *Marine Policy*, 36(2), 520–527. <https://doi.org/10.1016/j.marpol.2011.10.002>
- Ponte, S., Raakjær, J., & Campling, L. (2007). Swimming upstream: Market access for African fish exports in the context of WTO and EU negotiations and regulation. *Development Policy Review*, 25(1), 113–138. <https://doi.org/10.1111/j.1467-7679.2007.00362.x>
- Poos, M. S., & Jackson, D. A. (2011). Addressing the removal of rare species in multivariate bioassessments: The impact of methodological choices. *Ecological Indicators*, 18, 82–90.
- Postaire, B. D., Bakker, J., Gardiner, J., Wiley, T. R., & Chapman, D. D. (2020). Environmental DNA detection tracks established seasonal occurrence of blacktip sharks (*Carcharhinus limbatus*) in a semi-enclosed subtropical bay. *Scientific Reports*, 10(1), 1–8. <https://doi.org/10.1038/s41598-020-68843-0>
- Poulakis, G. R., Urakawa, H., Stevens, P. W., DeAngelo, J. A., Timmers, A. A., Grubbs, R. D., & Olin, J. A. (2017). Sympatric elasmobranchs and fecal samples provide insight into the trophic ecology of the smalltooth sawfish. *Endangered Species Research*, 32, 491–506. <https://doi.org/10.3354/ESR00824>
- Power, M. E., Tilman, D., Estes, J. A., Menge, B. A., Bond, W. J., Mills, L. S., & Paine, R. T. (1996). Challenges in the quest for keystones defining keystones. *BioScience*, 46(8), 609–620.
- Pradhan, S. K., Nayak, P. K., & Armitage, D. (2022). A social-ecological systems perspective on dried fish value chains. *Current Research in Environmental Sustainability*, 4(September 2021), 100128. <https://doi.org/10.1016/j.crsust.2022.100128>
- Prasetyo, A. P., McDevitt, A. D., Murray, J. M., Barry, J., Agung, F., Muttaqin, E., & Mariani, S. (2021). Shark and ray trade in and out of Indonesia: Addressing knowledge gaps on the path to sustainability. *Marine Policy*, 133(July), 104714. <https://doi.org/10.1016/j.marpol.2021.104714>
- Pridmore, R. D., Thrush, S. F., Hewitt, J. E., & Roper, D. S. (1990). Macrobenthic community composition of six intertidal sandflats in Manukau Harbour, New Zealand. *New Zealand Journal of Marine and Freshwater Research*, 24(1), 81–96. <https://doi.org/10.1080/00288330.1990.9516404>
- Prieto, G. (2023). Shark fisheries during the second millennium BC in Gramalote, north coast of Peru. *Journal of Island and Coastal Archaeology*, 18(2), 165–195. <https://doi.org/10.1080/15564894.2021.1910386>
- Puniwai, N. (2020). Pua ka wiliwili, nanahu ka manō: Understanding sharks in Hawaiian culture. *Human Biology*, 92(1), 11–17. <https://doi.org/10.13110/humanbiology.92.1.03>
- Punt, A. E., Pribac, F., Walker, T. I., Taylor, B. L., & Prince, J. D. (2000). Stock assessment of school shark, *Galeorhinus galeus*, based on a spatially explicit population dynamics model. *Marine and Freshwater Research*, 51(3), 205–220. <https://doi.org/10.1071/MF99124>
- Purvis, A., Gittleman, J. L., Cowlishaw, G., & Mace, G. M. (2000). Predicting extinction risk in declining species. *Proceedings of the Royal Society B: Biological Sciences*, 267(1456), 1947–1952. <https://doi.org/10.1098/rspb.2000.1234>

Pytka, J. M., Kyne, P. M., Carlson, J. K., Wosnick, N., & Jabado, R. W. (2023). A tangled web: Global review of fishing interactions with rhino rays. *Reviews in Fish Biology and Fisheries*, 0123456789. <https://doi.org/10.1007/s11160-023-09821-3>

Q

- QGIS Development Team & Others. (2018). QGIS. Open Source Geospatial Foundation Project.
- Queiroz, N., Humphries, N. E., Couto, A., Vedor, M., Costa, I., Sequeira, A. M. M., & Sims, D. W. (2019). Global spatial risk assessment of sharks under the footprint of fisheries. *Nature*, 572(7770), 461–466. <https://doi.org/10.1038/s41586-019-1444-4>
- Quezada-Romegialli, C., Jackson, A. L., Hayden, B., Kahilainen, K. K., Lopes, C., & Harrod, C. (2018). tRophicPosition, an R package for the Bayesian estimation of trophic position from consumer stable isotope ratios. *Methods in Ecology and Evolution*, 9(6), 1592–1599. <https://doi.org/10.1111/2041-210X.13009>

R

- R Core Team. (2017). R: A language and environment for statistical computing. <https://www.R-project.org/>
- Rafaelli, D., & Hawkins, S. J. (1999). *Intertidal Ecology* (1st ed.). Kluwer Academic Publishers. <https://doi.org/10.2307/1352485>
- RAMSAR. (2014). Archipel Bolama-Bijagós. RAMSAR. <https://rsis.ramsar.org/ris/2198>
- Ramsar Convention Secretariat. (2014). Ramsar Sites Information Service. RAMSAR. <https://rsis.ramsar.org/ris/2198>.
- Rangeley, R. W., & Kramer, D. L. (1995). Use of rocky intertidal habitats by juvenile pollock *Pollachius virens*. *Marine Ecology Progress Series*, 126(1–3), 9–17. <https://doi.org/10.3354/meps126009>
- Raoult, V., Broadhurst, M. K., Peddemors, V. M., Williamson, J. E., & Gaston, T. F. (2019). Resource use of great hammerhead sharks (*Sphyrna mokarran*) off eastern Australia. *Journal of Fish Biology*. <https://doi.org/10.1111/jfb.14160>
- Rasher, D. B., Hoey, A. S., & Hay, M. E. (2017). Cascading predator effects in a Fijian coral reef ecosystem. *Scientific Reports*, 7(1), 1–10. <https://doi.org/10.1038/s41598-017-15679-w>
- Raymond, W. W., Barber, J. S., Dethier, M. N., Hayford, H. A., Harley, C. D. G., King, T. L., & McDonald, P. S. (2022). Assessment of the impacts of an unprecedented heatwave on intertidal shellfish of the Salish Sea. *Ecology*, 1–7. <https://doi.org/10.1002/ecy.3798>
- Reid, W. D. K., Sweeting, C. J., Wigham, B. D., McGill, R. A. R., & Polunin, N. V. C. (2016). Isotopic niche variability in macroconsumers of the East Scotia Ridge (Southern Ocean) hydrothermal vents: What more can we learn from an ellipse? *Marine Ecology Progress Series*, 542, 13–24. <https://doi.org/10.3354/meps11571>
- Reidenauer, J. A., & Thistle, D. (1981). Response of a soft-bottom harpacticoid community to stingray (*Dasyatis sabinia*) disturbance. *Marine Biology*, 65(3), 261–267. <https://doi.org/10.1007/BF00397120>
- Renz, J. R., Powilleit, M., Gogina, M., Zettler, M. L., Morys, C., & Forster, S. (2018). Community bioirrigation potential (BIPc), an index to quantify the potential for solute exchange at the sediment-water interface. *Marine Environmental Research*, 141(September), 214–224. <https://doi.org/10.1016/j.marenvres.2018.09.013>

- Ricker, W. E. (1975). Computation and interpretation of biological statistics of fish populations. *Bulletin of the Fisheries Research Board of Canada*, 191, 1–382.
- Ries, J. B., Cohen, A. L., & McCorkle, D. C. (2009). Marine calcifiers exhibit mixed responses to CO₂-induced ocean acidification. *Geology*, 37(12), 1131–1134. <https://doi.org/10.1130/G30210A.1>
- Rigby, C. L., Dulvy, N. K., Barreto, R., Carlson, J., Fernando, D., Fordham, S., Francis, M. P., Herman, K., Jabado, R. W., Liu, K. M., Marshall, A., Pacoureau, N., Romanov, E., Sherley, R. B., & Winker, H. (2019). *Sphyrna lewini*. In *The IUCN red list of threatened species 2019*: E.T39385A2918526.
- Ripple, W. J., Estes, J. A., Beschta, R. L., Wilmers, C. C., Ritchie, E. G., Hebblewhite, M., & Wirsing, A. J. (2014). Status and ecological effects of the world's largest carnivores. *Science (New York, N.Y.)*, 343(6167). <https://doi.org/10.1126/science.1241484>
- Ritchie, E. G., & Johnson, C. N. (2009). Predator interactions, mesopredator release and biodiversity conservation. *Ecology Letters*, 12(9), 982–998. <https://doi.org/10.1111/j.1461-0248.2009.01347.x>
- Robillard, M., & Seret, B. (2006). Cultural importance and decline of sawfish (*Pristidae*) populations in West Africa. *Cybio*, 30(4), 23–30.
- Roff, G., Doropoulos, C., Rogers, A., Bozec, Y. M., Krueck, N. C., Aurellado, E., & Mumby, P. J. (2016). The ecological role of sharks on coral reefs. *Trends in Ecology and Evolution*, 31(5), 395–407. <https://doi.org/10.1016/j.tree.2016.02.014>
- Rose, M. D., Polis, G. A., Rose, M. D., & A, G. (2018). The distribution and abundance of coyotes: The effects of allochthonous food subsidies from the sea author. *Wiley on Behalf of the Ecological Society of America Stable URL*, 79(3), 998–1007.
- Rosenblatt, A. E., Heithaus, M. R., Mather, M. E., Matich, P., Nifong, J. C., Ripple, W. J., & Silliman, B. R. (2013). The roles of large top predators in coastal ecosystems new insights from long term ecological research. *Oceanography*, 26(3), 157–167. <https://doi.org/10.5670/oceanog.2013.59>
- Rosenfeld, J. S. (2002). Functional redundancy in ecology and conservation. *Oikos (Copenhagen, Denmark)*, 98(1), 156–162.
- Routley, M. H., Nilsson, G. E., & Renshaw, G. M. C. (2002). Exposure to hypoxia primes the respiratory and metabolic responses of the epaulette shark to progressive hypoxia. *Comparative Biochemistry and Physiology - A Molecular and Integrative Physiology*, 131(2), 313–321. [https://doi.org/10.1016/S1095-6433\(01\)00484-6](https://doi.org/10.1016/S1095-6433(01)00484-6)
- Ruiz, D. J., & Wolff, M. (2011). The Bolivar Channel Ecosystem of the Galapagos Marine Reserve: Energy flow structure and role of keystone groups. *Journal of Sea Research*, 66(2), 123–134. <https://doi.org/10.1016/j.seares.2011.05.006>
- Ruppert, J. L. W., Travers, M. J., Smith, L. L., Fortin, M.-J., & Meekan, M. G. (2013). Caught in the Middle: Combined Impacts of Shark Removal and Coral Loss on the Fish Communities of Coral Reefs. *PLoS One*, 8(9), 74648. <https://doi.org/10.1371/journal.pone.0074648>
- Russo, R. A. (2015). Observations of predation and loss among leopard sharks and brown smoothhounds in San Francisco Bay, California. *California Fish and Game*, 101(2), 149–157.

S

- Sáenz-Arroyo, A., Roberts, C. M., Torre, J., Cariño-Olvera, M., & Enríquez-Andrade, R. R. (2005). Rapidly shifting of among. *Proceedings: Biological Sciences*, 272(1575), 1957–1962.

- Salas, S., Chuenpagdee, R., Seijo, J. C., & Charles, A. (2007). Challenges in the assessment and management of small-scale fisheries in Latin America and the Caribbean. *Fisheries Research*, 87(1), 5–16. <https://doi.org/10.1016/j.fishres.2007.06.015>
- Salvig, J. C., Asbirk, S., Kjeldsen, J. P., & Rasmussen, P. A. F. (1994). Wintering waders in the Bijagós Archipelago, Guinea-Bissau 1992-1993. *Ardea*, 82(1), 137–142.
- Santo, V., & Bennett, W. A. (2011). Is post-feeding thermotaxis advantageous in elasmobranch fishes? *Journal of Fish Biology*, 78(1), 195–207. <https://doi.org/10.1111/j.1095-8649.2010.02853.x>
- Saucedo Barrón, C. J., Colado Uribe, G., Martínez Adrián, J. G., Burgos Zazueta, S., Chacón Cortez, J. G., & Espinoza Fierro, J. (1982). Contribución al estudio de la pesquería del tiburón en la zona sur de Sinaloa. Memoria de Servicio Social Universitario, Escuela de Ciencias del Mar, tesis Universidad Autónoma de Sinaloa.
- Schaffmeister, B. E., Hiddink, J. G., & Wolff, W. J. (2006). Habitat use of shrimps in the intertidal and shallow subtidal seagrass beds of the tropical Banc d'Arguin, Mauritania. *Journal of Sea Research*, 55(3), 230–243. <https://doi.org/10.1016/j.seares.2005.10.003>
- Scheiffarth, G., & Nehls, G. (1997). Consumption of benthic fauna by carnivorous birds in the Wadden Sea. *Helgolander Meeresuntersuchungen*, 51(3), 373–387. <https://doi.org/10.1007/BF02908721>
- Schlaff, A. M., Heupel, M. R., & Simpfendorfer, C. A. (2014). Influence of environmental factors on shark and ray movement, behaviour and habitat use: A review. *Reviews in Fish Biology and Fisheries*, 24(4), 1089–1103. <https://doi.org/10.1007/s11160-014-9364-8>
- Schmitz, O. J., Hawlena, D., & Trussell, G. C. (2010). Predator control of ecosystem nutrient dynamics. *Ecology Letters*, 13(10), 1199–1209. <https://doi.org/10.1111/j.1461-0248.2010.01511.x>
- Schweiss, K. E., Lehman, R. N., Drymon, J. M., & Phillips, N. M. (2020). Development of highly sensitive environmental DNA methods for the detection of Bull Sharks, *Carcharhinus leucas* (Müller and Henle, 1839). using Droplet Digital™ PCR. *Environmental DNA*, 2(1), 3–12. <https://doi.org/10.1002/edn3.39>
- Sciara, G. M. B., N., S., F., & Mancusi, C. M. (2015). *Dasyatis marmorata*, marbled stingray. In The IUCN red list of threatened species 2009 (p. 161748 5494614, 8235).
- Seguel, M. E., Sebastián Ramírez-Rivera, S., Concha, F., Espinoza, M., & Hernández, S. (2022). Predation and epibiosis on egg capsules of the Shorttail fanskate, *Sympterygia brevicaudata*. *Revista de Biología Marina y Oceanografía*, 57(Especial). <https://doi.org/10.22370/rbmo.2022.57.especial.3314>
- Seidu, I., Brobbey, L. K., Danquah, E., Oppong, S. K., Seidu, M., & Dulvy, N. K. (2021). Fishing for survival: Importance of shark fisheries for the livelihoods of coastal communities in Western Ghana. *Fisheries Research*, 246(October 2021), 1–42. <https://doi.org/10.1016/j.fishres.2021.106157>
- Séret, B., & Naylor, G. J. P. (2016). *Rhynchorhina mauritaniensis*, a new genus and species of wedgefish from the eastern central Atlantic (Elasmobranchii: Batoidea: Rhinidae). *Zootaxa*, 4138(2), 291–308. <https://doi.org/10.11646/zootaxa.4138.2.4>
- Sergio, F., Newton, I., Marchesi, L., & Pedrini, P. (2006). Ecologically justified charisma: Preservation of top predators delivers biodiversity conservation. *Journal of Applied Ecology*, 43(6), 1049–1055. <https://doi.org/10.1111/j.1365-2664.2006.01218.x>
- Sherman, C. S., Heupel, M. R., Moore, S. K., Chin, A., & Simpfendorfer, C. A. (2020). When sharks are away, rays will play: Effects of top predator removal in coral reef ecosystems. *Marine Ecology Progress Series*, 641, 145–157.

- Sherman, C. S., Simpfendorfer, C. A., Pacoureaux, N., Matsushiba, J. H., Yan, H. F., Walls, R. H. L., Rigby, C. L., VanderWright, W. J., Jabado, R. W., Pollom, R. A., Carlson, J. K., Charvet, P., Bin Ali, A., Fahmi, C., J. D., DH, H., KB, F., B, E., TD, P., ... N.K. (2023). Half a century of rising extinction risk of coral reef sharks and rays. *Nature Communications*, 14(15).
- Sherman, K. M., Reidenauer, J. A., Thistle, D., & Meeter, D. (1983). Role of a natural disturbance in an assemblage of marine free-living nematodes. *Marine Ecology Progress Series*, 11, 23–30.
- Shibuya, A., Souza Rosa, R., & Gadig, O. B. F. (2005). Stomach contents of *Galeocerdo cuvier* and *Carcharhinus plumbeus* (Elasmobranchii: Carcharhinidae) caught off Paraíba state, Brazil. *Arquivos de Ciências do Mar*, 38(1–2), 105–107.
- Shiffman, D., Bangley, C., & Macdonald, C. (2023). A prized Pacific shark: The rise and fall (and rise again...?) of the world's first ecolabel certified sustainable shark fishery. *Journal of Fish Biology*. <https://doi.org/10.1111/jfb.15467>
- Shiffman, D. S., & Hueter, R. E. (2017). A United States shark fin ban would undermine sustainable shark fisheries. *Marine Policy*, 85(July), 138–140. <https://doi.org/10.1016/j.marpol.2017.08.026>
- Shiple, O. N., Gallagher, A. J., Shiffman, D. S., Kaufman, L., & Hammerschlag, N. (2019). Diverse resource-use strategies in a large-bodied marine predator guild: Evidence from differential use of resource subsidies and intraspecific isotopic variation. *Marine Ecology Progress Series*, 623, 71–83. <https://doi.org/10.3354/meps12982>
- Sieben, K., Ljunggren, L., Bergström, U., & Eriksson, B. K. (2011). A meso-predator release of stickleback promotes recruitment of macroalgae in the Baltic Sea. *Journal of Experimental Marine Biology and Ecology*, 397(2), 79–84. <https://doi.org/10.1016/j.jembe.2010.11.020>
- Sievers, M., Brown, C. J., Tulloch, V. J. D., Pearson, R. M., Haig, J. A., Turschwell, M. P., & Connolly, R. M. (2019). The role of vegetated coastal wetlands for marine megafauna conservation. *Trends in Ecology and Evolution*, 34(9), 807–817. <https://doi.org/10.1016/j.tree.2019.04.004>
- Simpfendorfer, C. A., & Dulvy, N. K. (2017). Bright spots of sustainable shark fishing. *Current Biology*. <https://doi.org/10.1016/j.cub.2016.12.017>
- Simpfendorfer, C. A., Goodreid, A. B., & Mcauley, R. B. (2001). Size, sex and geographic variation in the diet of the tiger shark, *Galeocerdo cuvier*, from Western Australian waters. *Environmental Biology of Fishes*, 61(1), 37–46. <https://doi.org/10.1023/A:1011021710183>
- Simpfendorfer, C. A., & Milward, N. E. (1993). Utilisation of a tropical bay as a nursery area by sharks of the families Carcharhinidae and Sphyrnidae. *Environmental Biology of Fishes*, 37(4), 337–345. <https://doi.org/10.1007/BF00005200>
- Smale, M. J. (2005). The diet of the ragged-tooth shark *Carcharias taurus* Rafinesque 1810 in the Eastern Cape, South Africa. *African Journal of Marine Science*, 27(1), 331–335. <https://doi.org/10.2989/18142320509504091>
- Smart, J. J. (2016). Multimodel approaches in shark and ray growth studies: Strengths, weaknesses and the future. *Fish and Fisheries*, 17(4), 955–971. <https://doi.org/10.1111/faf.12154>
- Smith, J. W., & Merriner, J. V. (1985). Food habits and feeding behavior of the cownose ray, *Rhinoptera bonasus*, in lower Chesapeake Bay. *Estuaries*, 8(3), 305–310. <https://doi.org/10.2307/1351491>
- Smith, R. I. (1956). The ecology of the Tamar estuary. VII. Observations on the interstitial salinity of intertidal muds in the estuarine habitat of *Nereis diversicolor*. *Journal of the Marine Biological Association of the United Kingdom*, 35(1), 81–104. <https://doi.org/10.1017/S0025315400008985>

- Smith, S. E. (2005). Leopard shark mating observed off La Jolla, California. *California Fish and Game*, 91(2), 128–135.
- Smith, W. D., Cailliet, G. M., & Melendez, E. M. (2007). Maturity and growth characteristics of a commercially exploited stingray, *Dasyatis dipterura*. *Marine and Freshwater Research*, 58(1), 54–66. <https://doi.org/10.1071/MF06083>.
- Snelson, F. F., Mulligan, T. J., & Williams, S. E. (1984). Food habits, occurrence, and population structure of the bull shark, *Carcharhinus leucas*, in Florida coastal lagoons. *Bulletin of Marine Science*, 34(1), 71–80.
- Snelson, F. F., & Williams, S. E. (1981). Notes on the occurrence, distribution, and biology of elasmobranch fishes in the Indian River lagoon system, Florida. *Estuaries*, 4(2), 110–120. <https://doi.org/10.2307/1351673>
- Soto, C. G. (2006). Socio-cultural barriers to applying fisheries. In *Knowledge in fisheries management: An evaluation of literature cases thesis submitted in partial fulfilment of the requirements for the degree of doctor of philosophy in the school of resource and environmental*. Simon Fraser University.
- Spalding, M. D., Ruffo, S., Lacambra, C., Meliane, I., Hale, L. Z., Shepard, C. C., & Beck, M. W. (2014). The role of ecosystems in coastal protection: Adapting to climate change and coastal hazards. *Ocean and Coastal Management*, 90, 50–57. <https://doi.org/10.1016/j.ocecoaman.2013.09.007>
- Speed, C. W., Field, I. C., Meekan, M. G., & Bradshaw, C. J. A. (2010). Complexities of coastal shark movements and their implications for management. *Marine Ecology Progress Series*, 408, 275–293. <https://doi.org/10.3354/meps08581>
- Spiegelberger, T., & Ganslosser, U. (2005). Habitat analysis and exclusive bank feeding of the Antillean manatee (*Trichechus manatus manatus*) in the coswine swamps of French Guiana, South America. *Tropical Zoology*, 18(1), 1–12. <https://doi.org/10.1080/03946975.2005.10531210>
- Stål, J., Paulsen, S., Pihl, L., Rönnbäck, P., Söderqvist, T., & Wennhage, H. (2008). Coastal habitat support to fish and fisheries in Sweden: Integrating ecosystem functions into fisheries management. *Ocean and Coastal Management*, 51(8–9), 594–600. <https://doi.org/10.1016/j.ocecoaman.2008.06.006>
- Staples, D., & Funge-Smith, S. (2009). *Ecosystem approach to fisheries and aquaculture: Implementing the FAO Code of Conduct for Responsible Fisheries*. FAO Regional Office for Asia and the Pacific.
- Starr, R. M., Carr, M., Malone, D., Greenley, A., & McMillan, S. (2010). Complementary sampling methods to inform ecosystem-based management of nearshore fisheries. *Marine and Coastal Fisheries*, 2(1), 159–179. <https://doi.org/10.1577/c08-056.1>
- Stein, R. W., Mull, C. G., Kuhn, T. S., Aschliman, N. C., Davidson, L. N. K., Joy, J. B., & Mooers, A. O. (2018). Global priorities for conserving the evolutionary history of sharks, rays and chimaeras. *Nature Ecology & Evolution*, 2(2), 288–298. <https://doi.org/10.1038/s41559-017-0448-4>
- Stephens, D. W., Brown, J. S., & Ydenberg, R. C. (Eds.). (2007). *Foraging*. University of Chicago Press.
- Stevens, J. D. (1984). Life-history and ecology of sharks at Aldabra Atoll, Indian Ocean. *Proceedings of the Royal Society of London B: Biological Sciences*, 222(1226), 79–106.
- Stevens, J. D., Bonfil, R., Dulvy, N. K., & Walker, P. A. (2000). The effects of fishing on sharks, rays, and chimaeras (chondrichthyans), and the implications for marine ecosystems. *ICES Journal of Marine Science*, 57(3), 476–494. <https://doi.org/10.1006/jmsc.2000.0724>

- Stevens, J. D., & Lyle, J. M. (1989). Biology of three hammerhead sharks (*Eusphyrna blochii*, *Sphyrna mokarran* and *S. lewini*) from Northern Australia. *Marine and Freshwater Research*, 40(2), 46–129. <https://doi.org/10.1071/MF9890129>
- Stewart, K. A. (2019). Understanding the effects of biotic and abiotic factors on sources of aquatic environmental DNA. *Biodiversity and Conservation*, 28(5), 983–1001. <https://doi.org/10.1007/s10531-019-01709-8>
- Stillwell, C. E., & Kohler, N. E. (1982). Food, feeding habits, and estimates of daily ration of the shortfin mako (*Isurus oxyrinchus*) in the Northwest Atlantic. *Canadian Journal of Fisheries and Aquatic Sciences*, 39(3), 407–414. <https://doi.org/10.1139/f82-058>
- Stobart, B., Warwick, R., González, C., Mallol, S., Díaz, D., Reñones, O., & Goñi, R. (2009). Long-term and spillover effects of a marine protected area on an exploited fish community. *Marine Ecology Progress Series*, 384, 47–60. <https://doi.org/10.3354/meps08007>
- Stock, B. C., Jackson, A. L., Ward, E. J., Parnell, A. C., Phillips, D. L., & Semmens, B. X. (2018). Analyzing mixing systems using a new generation of Bayesian tracer mixing models. *PeerJ*, 2018(6), 1–27. <https://doi.org/10.7717/peerj.5096>
- Strickland, T. L. (2009). Critical habitat for the endangered distinct population segment of the smalltooth sawfish. *Federal Register*, 74(169), 45353–45378.
- Strong, W. R., Snelson, F. F., & Gruber, S. H. (1990). Hammerhead shark predation on stingrays: An observation of prey handling by *Sphyrna mokarran*. *Copeia*, 1990(3), 836–840.
- Stroud, D. A., Baker, A., Blanco, D. E., Davidson, N. C., Delany, S., Ganter, B., & Zöckler, C. (2006). The conservation and population status of the world's waders at the turn of the millennium. *Waterbirds around the World*, 643–648.
- Stuart-Smith, R. D., Bates, A. E., Lefcheck, J. S., Duffy, J. E., Baker, S. C., Thomson, R. J., & Edgar, G. J. (2013). Integrating abundance and functional traits reveals new global hotspots of fish diversity. *Nature*, 501(7468), 539–542. <https://doi.org/10.1038/nature12529>
- Studds, C. E., Kendall, B. E., Murray, N. J., Wilson, H. B., Rogers, D. I., Clemens, R. S., & Fuller, R. A. (2017). Rapid population decline in migratory shorebirds relying on Yellow Sea tidal mudflats as stopover sites. *Nature Communications*, 8, 1–7. <https://doi.org/10.1038/ncomms14895>
- Suchanek, T. H., & Colin, P. L. (1986). Rates and effects of bioturbation by invertebrates and fishes at Enewetak and Bikini Atolls. *Bulletin of Marine Science*, 38, 25–34.
- Summers, A. P. (2000). Stiffening the stingray skeleton—An investigation of durophagy in myliobatid stingrays (Chondrichthyes, Batoidea, Myliobatidae). *Journal of Morphology*, 243(2), 113–126. [https://doi.org/10.1002/\(SICI\)1097-4687\(200002\)243:2](https://doi.org/10.1002/(SICI)1097-4687(200002)243:2)
- Sutherland, W. J. (1982). Do oystercatchers select the most profitable cockles? *Animal Behaviour*, 30(3), 857–861. [https://doi.org/10.1016/S0003-3472\(82\)80159-0](https://doi.org/10.1016/S0003-3472(82)80159-0)
- Suykerbuyk, W., Bouma, T. J., Govers, L. L., Giesen, K., Jong, D. J., Herman, P., & Katwijk, M. M. (2016). Surviving in changing seascapes: Sediment dynamics as bottleneck for long-term seagrass presence. *Ecosystems (New York, N.Y.)*, 19(2), 296–310. <https://doi.org/10.1007/s10021-015-9932-3>
- Swanson, H. K., Lysy, M., Power, M., Stasko, A. D., Johnson, J. D., & Reist, J. (2015). A new probabilistic method for quantifying n-dimensional ecological niches and niche overlap. *Ecology*, 96(2), 318–324.
- Swartz, W., Sala, E., Tracey, S., Watson, R., & Pauly, D. (2010). The spatial expansion and ecological footprint of fisheries (1950 to present). *PLoS ONE*, 5(12), 3–8. <https://doi.org/10.1371/journal.pone.0015143>

T

- Taberlet, P., Bonin, A., Zinger, L., & Coissac, E. (2018). *Environmental DNA: For biodiversity research and monitoring*. Oxford University Press. <https://doi.org/10.1093/oso/9780198767220.001.0001>
- Takeuchi, S., Takahara, Y., Agata, Y., Nasuda, J., Yamada, F., & Tamaki, A. (2013). Response of suspension-feeding clams to natural removal of bioturbating shrimp on a large estuarine intertidal sandflat in western Kyushu, Japan. *Journal of Experimental Marine Biology and Ecology*, 448, 308–320. <https://doi.org/10.1016/j.jembe.2013.07.018>
- Takeuchi, S., & Tamaki, A. (2014). Assessment of benthic disturbance associated with stingray foraging for ghost shrimp by aerial survey over an intertidal sandflat. *Continental Shelf Research*, 84, 139–157. <https://doi.org/10.1016/j.csr.2014.05.007>
- Talent, L. G. (1982). Food habits of the gray smoothhound, *Mustelus californicus*, the brown smoothhound, *Mustelus henlei*, the shovelnose guitarfish, *Rhinobatos productus*, and the bat ray, *Miliobatis californica*, in Elkhorn Slough, California. *California Fish & Game*, 68(4), 224–234.
- Talwar, B. S., Bradley, D., Berry, C., Bond, M. E., Bouyoucos, I. A., Brooks, A. M. L., & Brooks, E. J. (2022). Estimated life-history traits and movements of the Caribbean reef shark (*Carcharhinus perezi*) in The Bahamas based on tag-recapture data. *Marine Biology*, 169(5), 1–19. <https://doi.org/10.1007/s00227-022-04044-9>
- Techera, E. J., & Klein, N. (2011). Fragmented governance: Reconciling legal strategies for shark conservation. *Marine Policy*, 35(1), 73–78. <https://doi.org/10.1016/j.marpol.2010.08.003>
- Teh, L. C. L., & Pauly, D. (2018). Who brings in the fish? The relative contribution of small-scale and industrial fisheries to food security in Southeast Asia. *Frontiers in Marine Science*, 4(FEB), 1–9. <https://doi.org/10.3389/fmars.2018.00044>
- Temmerman, S., Meire, P., Bouma, T. J., Herman, P. M. J., Ysebaert, T., & Vriend, H. J. (2013). Ecosystem-based coastal defence in the face of global change. *Nature*, 504, 79–83.
- Temple, A. J. (2020). Life-history, exploitation and extinction risk of the data-poor Baraka's whipray (*Maculabatis ambigua*) in small-scale tropical fisheries. *Journal of Fish Biology*, 97(3), 708–719. <https://doi.org/10.1111/jfb.14425>
- Temple, A. J., Wambiji, N., Poonian, C. N. S., Jiddawi, N., Stead, S. M., Kiszka, J. J., & Berggren, P. (2019). Marine megafauna catch in southwestern Indian Ocean small-scale fisheries from landings data. *Biological Conservation*, 230(August 2018), 113–121. <https://doi.org/10.1016/j.biocon.2018.12.024>
- Ten key biological traits of marine benthic invertebrates surveyed in Northwest Europe (Version 2). (2022). [dataset]. Cefas. <https://doi.org/10.14466/CEFASDATAHUB.123>
- Terborgh, J., & Estes, J. A. (Eds.). (2010). *Trophic cascades: Predators, prey, and the changing dynamics of nature* (Vol. 23). Island Press. <https://doi.org/10.1007/s12045-018-0730-z>
- Tesfamichael, D., Pitcher, T. J., & Pauly, D. (2014). Assessing changes in fisheries using fishers' knowledge to generate long time series of catch rates: A case study from the red sea. *Ecology and Society*, 19(1). <https://doi.org/10.5751/ES-06151-190118>
- Tessier, A., Descloux, S., Lae, R., Cottet, M., Guedant, P., & Guillard, J. (2016). Fish assemblages in large tropical reservoirs: Overview of fish population monitoring methods. *Reviews in Fisheries Science and Aquaculture*, 24(2), 160–177. <https://doi.org/10.1080/23308249.2015.1112766>

- Tetley, M. J., Braulik, G. T., Lanfredi, C., Minton, G., Panigada, S., Politi, E., & Hoyt, E. (2022). The important marine mammal area network: A tool for systematic spatial planning in response to the marine mammal habitat conservation crisis. *Frontiers in Marine Science*, 9(March), 1–13. <https://doi.org/10.3389/fmars.2022.841789>
- Thaman, R. R., Puia, T., Tongabaea, W., Namona, A., & Fong, T. (2010). Marine biodiversity and ethnobiodiversity of Bellona (Mungiki) Island, Solomon Islands. *Singapore Journal of Tropical Geography*, 31(1), 70–84. <https://doi.org/10.1111/j.1467-9493.2010.00391.x>
- Thiault, L., Collin, A., Chlous, F., Gelcich, S., & Claudet, J. (2017). Combining participatory and socioeconomic approaches to map fishing effort in smallscale fisheries. *PLoS ONE*, 12(5), 1–18. <https://doi.org/10.1371/journal.pone.0176862>
- Thistle, D. (1981). Natural physical disturbances and communities of marine soft bottoms. *Marine Ecology Progress Series*, 6, 223–228.
- Thomsen, P. F., Kielgast, J., Iversen, L. L., Møller, P. R., Rasmussen, M., & Willerslev, E. (2012). Detection of a diverse marine fish fauna using environmental DNA from seawater samples. *PLoS ONE*, 7(8), 1–9. <https://doi.org/10.1371/journal.pone.0041732>
- Thorrold, S. R., Afonso, P., Fontes, J., Braun, C. D., Santos, R. S., Skomal, G. B., & Berumen, M. L. (2014). Extreme diving behaviour in devil rays links surface waters and the deep ocean. *Nature Communications*, 5, 1–7. <https://doi.org/10.1038/ncomms5274>
- Thrush, S. F., Hewitt, J. E., Gibbs, M., Lundquist, C., & Norkko, A. (2006). Functional role of large organisms in intertidal communities: Community effects and ecosystem function. *Ecosystems (New York, N.Y.)*, 9(6), 1029–1040. <https://doi.org/10.1007/s10021-005-0068-8>
- Thrush, S. F., Pridmore, R. D., Hewitt, J. E., & Cummings, V. J. (1994). The importance of predators on a sandflat—Interplay between seasonal changes in prey densities and predator effects. *Marine Ecology Progress Series*, 107(3), 211–222. <https://doi.org/10.3354/meps107211>
- Tillett, B. J., Meekan, M. G., & Field, I. C. (2014). Dietary overlap and partitioning among three sympatric carcharhinid sharks. *Endangered Species Research*, 25(3), 283–293. <https://doi.org/10.3354/esr00615>
- TinHan, T. C., & Wells, R. J. D. (2021). Spatial and ontogenetic patterns in the trophic ecology of juvenile bull sharks (*Carcharhinus leucas*) from the northwest Gulf of Mexico. *Frontiers in Marine Science*, 8(July), 1–12. <https://doi.org/10.3389/fmars.2021.664316>
- Tinker, M. T., Bentall, G., & Estes, J. A. (2008). Food limitation leads to behavioral diversification and dietary specialization in sea otters. *Proceedings of the National Academy of Sciences of the United States of America*, 105(2), 560–565. <https://doi.org/10.1073/pnas.0709263105>
- Tittensor, D. P., Mora, C., Jetz, W., Lotze, H. K., Ricard, D., Berghe, E. V., & Worm, B. (2010). Global patterns and predictors of marine biodiversity across taxa. *Nature*, 466(7310), 1098–1101. <https://doi.org/10.1038/nature09329>
- Tobin, A. J., Mapleston, A., Harry, A. V., & Espinoza, M. (2014). Big fish in shallow water; use of an intertidal surf-zone habitat by large-bodied teleosts and elasmobranchs in tropical northern Australia. *Environmental Biology of Fishes*, 97(7), 821–838. <https://doi.org/10.1007/s10641-013-0182-y>
- Todd, P. A., Ong, X., & Chou, L. M. (2010). Impacts of pollution on marine life in Southeast Asia. *Biodiversity and Conservation*, 19(4), 1063–1082. <https://doi.org/10.1007/s10531-010-9778-0>

- Tojo, H. (1996). Habitat selection, foraging behaviour and prey of five heron species in Japan. *Japanese Journal of Ornithology*, 45(3), 141–158. <https://doi.org/10.3838/jjo.45.141>
- Tréguier, A., Paillisson, J. M., Dejean, T., Valentini, A., Schlaepfer, M. A., & Roussel, J. M. (2014). Environmental DNA surveillance for invertebrate species: Advantages and technical limitations to detect invasive crayfish *Procambarus clarkii* in freshwater ponds. *Journal of Applied Ecology*, 51(4), 871–879. <https://doi.org/10.1111/1365-2664.12262>
- Turner, T. F., Collyer, M. L., & Krabbenhoft, T. J. (2010). A general hypothesis-testing framework for stable isotope ratios in ecological studies. *Ecology*, 91(8), 2227–2233. <https://doi.org/10.1890/09-1454.1>

U

- UNCTAD. (2020). The COVID-19 Pandemic and the Blue Economy: New challenges and prospects for recovery and resilience.
- UNEP-WCMC. (2019). Protected planet: Marine protected areas.
- UNESCO. (2020). Boloma-Bijagós Biosphere Reserve. UNESCO.

V

- Valadou, B., Brethes, J., & Inejih, C. (2006). Observations biologiques sur cinq espèces d'Élasmobranches du Parc National du Banc d'Arguin (Mauritanie). *Cybium*, 30(4), 313–322.
- Van Waerebeek, K., & Read, A. J. (1994). Reproduction of Dusky Dolphins, *Lagenorhynchus obscurus*, from Coastal Peru. *Journal of Mammalogy*, 75(4), 1054–1062. <https://doi.org/10.2307/1382489>
- VanBlaricom, G. R. (1982). Experimental analyses of structural regulation in a marine sand community exposed to oceanic swell. *Ecological Monographs*, 52(3), 283–305. <https://doi.org/10.2307/2937332>
- Vandermeer, J., & Pascual, M. (2006). Competitive coexistence through intermediate polyphagy. *Ecological Complexity*, 3(1), 37–43. <https://doi.org/10.1016/j.ecocom.2005.05.005>
- Vaudo, J. J., & Heithaus, M. R. (2011). Dietary niche overlap in a nearshore elasmobranch mesopredator community. *Marine Ecology Progress Series*, 425, 247–260. <https://doi.org/10.3354/meps08988>.
- Vaudo, J. J., & Heithaus, M. R. (2012). Diel and seasonal variation in the use of a nearshore sandflat by a ray community in a near pristine system. *Marine and Freshwater Research*, 63(11), 1077–1084. <https://doi.org/10.1071/MF11226>
- Vehtari, A., Gelman, A., & Gabry, J. (2017). Practical Bayesian model evaluation using leave-one-out cross-validation and WAIC. *Statistics and Computing*, 27(5), 1413–1432. <https://doi.org/10.1007/s11222-016-9696-4>.
- Venter, O., Magrath, A., Outram, N., Klein, C. J., Possingham, H. P., Marco, M., & Watson, J. E. M. (2018). Bias in protected-area location and its effects on long-term aspirations of biodiversity conventions. *Conservation Biology*, 32(1), 127–134. <https://doi.org/10.1111/cobi.12970>
- Veríssimo, A., Sampaio, Í., McDowell, J. R., Alexandrino, P., Mucientes, G., Queiroz, N., & Noble, L. R. (2017). World without borders—Genetic population structure of a highly migratory marine predator, the blue shark (*Prionace glauca*). *Ecology and Evolution*, 7(13), 4768–4781. <https://doi.org/10.1002/ece3.2987>
- Vermeulen, E. (2018). Intertidal habitat use of bottlenose dolphins (*Tursiops truncatus*) in Bahía Aan Antonio, Argentina. *Journal of the Marine Biological Association of the United Kingdom*, 98(5), 1109–1118. <https://doi.org/10.1017/S0025315417000856>

- Vianna, G. M. S., Hehre, E. J., White, R., Hood, L., Derrick, B., & Zeller, D. (2020). Long-term fishing catch and effort trends in the republic of the marshall islands, with emphasis on the small-scale sectors. *Frontiers in Marine Science*, 6(January), 1–13. <https://doi.org/10.3389/fmars.2019.00828>
- Virnstein, R. W. (1977). The importance of predation by crabs and fishes on benthic infauna in Chesapeake Bay. *Ecology*, 58(6), 1199–1217. <https://doi.org/10.2307/1935076>
- Visser, I. (1999). Benthic foraging on stingrays by killer whales (*Orcinus orca*) in New Zealand waters. *Marine Mammal Science*, 15(1), 220–227.
- Vlaming, V. L., & Sage, M. (1973). Osmoregulation in the euryhaline elasmobranch, *Dasyatis sabina*. *Comparative Biochemistry and Physiology – Part A: Physiology*, 45(1), 31–44. [https://doi.org/10.1016/0300-9629\(73\)90006-6](https://doi.org/10.1016/0300-9629(73)90006-6)
- Volpov, B. L., Hoskins, A. J., Battaile, B. C., Viviant, M., Wheatley, K. E., Marshall, G., & Arnould, J. P. Y. (2015). Identification of prey captures in Australian fur seals (*Arctocephalus pusillus doriferus*) using head-mounted accelerometers: Field validation with animal-borne video cameras. *PLoS ONE*, 10(6), 1–19. <https://doi.org/10.1371/journal.pone.0128789>

W

- Walker, P., & Hislop, J. (1998). Sensitive skates or resilient rays? Spatial and temporal shifts in ray species composition in the central and north-western North Sea between 1930 and the present day. *ICES Journal of Marine Science*, 55, 392–402. <https://doi.org/10.1006/jmsc.1997.0325>
- Wambiji, N., Kadagi, N. I., Everett, B. I., Temple, A. J., Kiszka, J. J., Kimani, E., & Berggren, P. (2022). Integrating long-term citizen science data and contemporary artisanal fishery survey data to investigate recreational and small-scale shark fisheries in Kenya. *Aquatic Conservation: Marine and Freshwater Ecosystems*, 32(8), 1306–1322. <https://doi.org/10.1002/aqc.3829>
- Wang, S., Brose, U., & Gravel, D. (2019). Intraguild predation enhances biodiversity and functioning in complex food webs. *Ecology*, 100(3), 1–10. <https://doi.org/10.1002/ecy.2616>
- Wang, Y., Wang, Y. P., Yu, Q., Du, Z., Wang, Z. B., & Gao, S. (2019). Sand–mud tidal flat morphodynamics influenced by alongshore tidal currents. *Journal of geophysical research*. *Oceans*, 124, 3818–3836.
- Wanink, J. H., & Zwarts, L. (1993). Environmental effects of the growth rate of intertidal invertebrates and some implications for foraging waders. *Netherlands Journal of Sea Research*, 31(4), 407–418. [https://doi.org/10.1016/0077-7579\(93\)90057-Y](https://doi.org/10.1016/0077-7579(93)90057-Y)
- Ward, P., & Myers, R. A. (2005). Shifts in open-ocean fish communities coinciding with the commencement of commercial fishing. *Ecology*, 86(4), 835–847. <https://doi.org/10.1890/03-0746>
- Watson, G. J., Murray, J. M., Schaefer, M., & Bonner, A. (2017). Bait worms: A valuable and important fishery with implications for fisheries and conservation management. *Fish and Fisheries*, 18(2), 374–388. <https://doi.org/10.1111/faf.12178>
- Watson, J. E. M., Dudley, N., Segan, D. B., & Hockings, M. (2014). The performance and potential of protected areas. *Nature*, 515(7525), 67–73. <https://doi.org/10.1038/nature13947>
- Weichselbaum, E., Costa, H., & Benelam, B. (2009). Synthesis report No 6: Traditional Foods in Europe. EUROFIR.

- Welch, H., Clavelle, T., White, T. D., Cimino, M. A., Osdel, J., Hochberg, T., & Hazen, E. L. (2022). Hot spots of unseen fishing vessels. *Science Advances*, 8(44), 1–11. <https://doi.org/10.1126/sciadv.abq2109>
- Wheatly, M. G. (1988). Integrated responses to salinity fluctuation. *American Zoologist*, 28(1), 65–77.
- White, A. T., Vogt, H. P., & Arin, T. (2000). Philippine coral reefs under threat: The economic losses caused by reef destruction. *Marine Pollution Bulletin*, 40(7), 598–605. [https://doi.org/10.1016/S0025-326X\(00\)00022-9](https://doi.org/10.1016/S0025-326X(00)00022-9)
- White, J., Heupel, M. R., Simpfendorfer, C. A., & Tobin, A. J. (2013). Shark-like batoids in pacific fisheries: Prevalence and conservation concerns. *Endangered Species Research*, 19(3), 277–284. <https://doi.org/10.3354/esr00473>
- White, T. D., Carlisle, A. B., Kroodsmas, D. A., Block, B. A., Casagrandi, R., Leo, G. A., & McCauley, D. J. (2017). Assessing the effectiveness of a large marine protected area for reef shark conservation. *Biological Conservation*, 207, 64–71. <https://doi.org/10.1016/j.biocon.2017.01.009>
- White, W. T., O'Neill, H., & Naylor, G. P. (2022). Taxonomy and diversity of extant elasmobranchs. In J. C. Carrier, C. A. Simpfendorfer, M. R. Heithaus, & K. E. Yopak (Eds.), *Biology of sharks and their relatives* (third (pp. 31–58). CRC Press.
- White, W. T., & Potter, I. C. (2004). Habitat partitioning among four elasmobranch species in nearshore, shallow waters of a subtropical embayment in Western Australia. *Marine Biology*, 145(5), 1023–1032. <https://doi.org/10.1007/s00227-004-1386-7>
- Williams, J. J., Papastamatiou, Y. P., Caselle, J. E., Bradley, D., & Jacoby, D. M. P. (2018). Mobile marine predators: An understudied source of nutrients to coral reefs in an unfished atoll. *Proceedings of the Royal Society B: Biological Sciences*, 285(1875). <https://doi.org/10.1098/rspb.2017.2456>
- Williamson, M. J., Tebbs, E. J., Dawson, T. P., & Jacoby, D. M. P. (2019). Satellite Remote Sensing in Shark and Ray Ecology, Conservation and Management. *Frontiers in Marine Science*, 6, 135. <https://doi.org/10.3389/fmars.2019.00135>
- Wilson, S. C., & Jones, K. A. (2018). Behaviour of harbour seal (*Phoca vitulina vitulina*) mother-pup pairs in irish sea intertidal habitats. *Biology and Environment*, 118B(1), 13–27. <https://doi.org/10.3318/BIOE.2018.02>
- Winemiller, K. O., & Rose, K. A. (1992). Patterns of life-history diversification in North American fishes: Implications for population regulation. *Canadian Journal of Fisheries and Aquatic Sciences*, 49, 2196–2218.
- Wirsing, A. J., Heithaus, M. R., & Dill, L. M. (2007). Fear factor: Do dugongs (*Dugong dugon*) trade food for safety from tiger sharks (*Galeocerdo cuvier*)? *Oecologia*, 153(4), 1031–1040. <https://doi.org/10.1007/s00442-007-0802-3>
- Wise, G., Mulvey, J. M., & Renshaw, G. M. C. (1998). Hypoxia tolerance in the epaulette shark (*Hemiscyllium ocellatum*). *Journal of Experimental Zoology*, 281(1), 1–5. [https://doi.org/10.1002/\(SICI\)1097-010X\(19980501\)281:13.0.CO;2-S](https://doi.org/10.1002/(SICI)1097-010X(19980501)281:13.0.CO;2-S)
- Wolff, W. J. (2005). The exploitation of living resources in the Dutch Wadden Sea: A historical overview. *Helgoland Marine Research*, 59(1), 31–38. <https://doi.org/10.1007/s10152-004-0204-4>
- Wood, S. N. (2004). Stable and Efficient Multiple Smoothing Parameter Estimation for Generalized Additive Models. *Journal of the American Statistical Association*, 99(467), 673–686. <https://doi.org/10.1198/016214504000000980>
- Wood, S. N. (2017). *Generalized additive models: An introduction with R* (second edition. 2nd Editio. Chapman an). <https://reseau-mexico.fr/sites/reseau-mexico.fr/files/igam.pdf>

- Worm, B., Barbier, E. B., Beaumont, N., Duffy, J. E., Folke, C., Halpern, B. S., & Watson, R. (2006). Impacts of biodiversity loss on ocean ecosystem services. *Science (New York, N.Y.)*, 314(5800), 787–790. <https://doi.org/10.1126/science.1132294>
- Worm, B., Davis, B., Kettener, L., Ward-Paige, C. A., Chapman, D., Heithaus, M. R., & Gruber, S. H. (2013). Global catches, exploitation rates, and rebuilding options for sharks. *Marine Policy*, 40(1), 194–204. <https://doi.org/10.1016/j.marpol.2012.12.034>
- Worm, B., Hilborn, R., Baum, J. K., Branch, T. A., Collie, J. S., Costello, C., & Zeller, D. (2009). Rebuilding Global Fisheries. *Science (New York, N.Y.)*, 325(5940), 578–585. <https://doi.org/10.1126/science.1173146>
- Wosnick, N., Leite, R. D., Giaretta, E. P., Morick, D., & Musyl, M. (2022). Global assessment of shark strandings. *Fish and Fisheries*, 23(4), 786–799. <https://doi.org/10.1111/faf.12648>
- Wosnick, N., Rios, R. M., Rincon, G., & Nunes, J. L. S. (2019). Evidence of tide pool use by a viviparous elasmobranch as a parturition site. *Acta Ichthyologica et Piscatoria*, 49(2), 189–193. <https://doi.org/10.3750/AIEP/02546>

Y

- Yadav, V. K., Jahageerdar, S., & Adinarayana, J. (2020). Use of different modeling approach for sensitivity analysis in predicting the Catch per Unit Effort (CPUE) of fish. *Indian J. Geo-Marine Sci*, 49, 129–1741.
- Yan, H. F., Kyne, P. M., Jabado, R. W., Leeney, R. H., Davidson, L. N. K., Derrick, D. H., & Dulvy, N. K. (2021). Overfishing and habitat loss drives range contraction of iconic marine fishes to near extinction. *Science Advances*, 7(7), 1–11. <https://doi.org/10.1126/sciadv.abb6026>
- Yao, Y. (2018). Using stacking to average bayesian predictive distributions. *Bayesian Analysis*, 13(3), 917–1007. <https://doi.org/10.1214/17-BA1091>.
- Yates, P. M., Tobin, A. J., Heupel, M. R., & Simpfendorfer, C. A. (2016). Benefits of marine protected areas for tropical coastal sharks. *Aquatic Conservation: Marine and Freshwater Ecosystems*, 26(6), 1063–1080. <https://doi.org/10.1002/aqc.2616>
- Yick, J. L., Tracey, S. R., & White, R. W. G. (2011). Niche overlap and trophic resource partitioning of two sympatric batoids co-inhabiting an estuarine system in southeast Australia. *Journal of Applied Ichthyology*, 27(5), 1272–1277. <https://doi.org/10.1111/j.1439-0426.2011.01819.x>
- Yigin, C. C., & Ismen, A. (2012). Age, growth and reproduction of the common stingray, *Dasyatis pastinaca* from the North Aegean Sea. *Marine Biology Research*, 8(7), 644–653. <https://doi.org/10.1080/17451000.2012.659667>

Z

- Zajac, R. N., Lewis, R. S., Poppe, L. J., Twichell, D. C., Vozarik, J., & DiGiacomo-Cohen, M. L. (2003). Responses of infaunal populations to benthoscape structure and the potential importance of transition zones. *Limnology and Oceanography*, 48(2), 829–842. <https://doi.org/10.4319/lo.2003.48.2.0829>
- Zanella, I., López-Garro, A., & Cure, K. (2019). Golfo Dulce: Critical habitat and nursery area for juvenile scalloped hammerhead sharks *Sphyrna lewini* in the Eastern Tropical Pacific Seascape. *Environmental Biology of Fishes*, 102(10), 1291–1300. <https://doi.org/10.1007/s10641-019-00907-1>
- Zaret, T. M., & Rand, A. S. (1971). Competition in tropical stream fishes: Support for the competitive exclusion principle. *Ecology*, 52(2), 336–342. <https://doi.org/10.2307/1934593>

References

- Zee, M., Angelini C, Govers LL, Christianen MJA, Altieri AH, Reijden KJ, Silliman BR, Koppel J, Geest M, Gils JA, Veer HW, Piersma T, Ruiter PC, Olff H, & Heide T. (2016). How habitat-modifying organisms structure the food web of two coastal ecosystems. *Proceedings of the Royal Society B: Biological Sciences* 283. <https://doi.org/10.1098/rspb.2015.2326>
- Zeeberg, J. J., Corten, A., & Graaf, E. (2006). Bycatch and release of pelagic megafauna in industrial trawler fisheries off Northwest Africa. *Fisheries Research*, 78(2-3), 186-195. <https://doi.org/10.1016/j.fishres.2006.01.012>
- Zharikov, Y., & Skilleter, G. A. (2003). Depletion of benthic invertebrates by bar-tailed godwits *Limosa lapponica* in a subtropical estuary. *Marine Ecology Progress Series*, 254, 151-162. <https://doi.org/10.3354/meps254151>
- Zwarts, L., & Blomert, A. M. (1992). Why knot *Calidris canutus* take medium-sized *Macoma balthica* when six prey species are available. *Marine Ecology Progress Series*, 83(2-3), 113-128. <https://doi.org/10.3354/meps083113>
- Zwarts, L., & Ens, B. J. (1999). Predation by birds on marine tidal flats. *International Ornithological Congress*, Durban, South Africa.
- Zwarts, L., & Piersma, T. (1990). How important is the Banc d'Arguin, Mauritania, as a temporary staging area for waders in spring? *Ardea*, 78(1-2), 113-121.
- Zwarts, L., & Wanink, J. (1984). How oystercatchers and curlews successively deplete clams. In *Coastal waders and wildfowl in winter* (pp. 69-83).
- Zwarts, L., & Wanink, J. (1989). Siphon size and burying depth in deposit- and suspension-feeding benthic bivalves. *Marine Biology*, 100(2), 227-240. <https://doi.org/10.1007/BF00391963>
- Zwarts, L., & Wanink, J. H. (1993). How the food supply harvestable by waders in the Wadden Sea depends on the variation in energy density, body weight, biomass, burying depth and behaviour of tidal-flat invertebrates. *Netherlands Journal of Sea Research*, 31(4), 441-476. [https://doi.org/10.1016/0077-7579\(93\)90059-2](https://doi.org/10.1016/0077-7579(93)90059-2)